

Two-body problem in Scalar-Tensor theories, an Effective-One-Body approach

Félix-Louis Julié

GARYFEST: Gravitation, Solitons and Symmetries

March 23rd 2017

[arXiv:1703.05360](https://arxiv.org/abs/1703.05360) FLJ - Nathalie Deruelle

- [GW150914](#) : The very first observation of a BBH coalescence by LIGO-Virgo has opened **a new era in gravitational wave astronomy**.
- Opportunity to bring **new tests of modified gravities**, in the strong-field regime near merger, a topic which is for the moment still in infancy.
- The “Effective-One-Body” (EOB) approach has proven to be a very powerful way to describe **analytically** the coalescence of 2 compact objects in **General Relativity**, from inspiral to merger.

Our proposition [arXiv:1703.05360]

- Can we extend the EOB approach to modified gravities ?
 - Consider the simplest and most studied example of **massless Scalar-Tensor theories**.
-
- First building block : map the conservative part of the two-body dynamics onto the geodesic of an effective metric.
 - ST-extension of [Buonanno-Damour 98]

We adopt the conventions of Damour and Esposito-Farèse [DEF 92, 95]

ST action in the Einstein-frame ($G_* \equiv c \equiv 1$)

$$S_{EF} = \frac{1}{16\pi} \int d^4x \sqrt{-g} \left(R - 2g^{\mu\nu} \partial_\mu \varphi \partial_\nu \varphi \right) + S_m \left[\Psi, \mathcal{A}^2(\varphi) g_{\mu\nu} \right]$$

- **Einstein metric** $g_{\mu\nu}$ free dynamics : Einstein-Hilbert term ; ordinary kinematical term for φ ;
- **BUT** matter Ψ is minimally coupled to the **Jordan metric** $\tilde{g}_{\mu\nu}$:

$$\tilde{g}_{\mu\nu} \equiv \mathcal{A}^2(\varphi) g_{\mu\nu}$$

where $\mathcal{A}(\varphi)$ **defines** the ST theory (GR : $\mathcal{A}(\varphi) = cst$).

- Encompass the Einstein Equivalence Principle

what about S_m ?

N-body problem in Scalar-Tensor theories

Phenomenological approach : Skeletonize extended bodies as point particles

- Non-self-gravitating case

$$S_m = - \sum_A \int d\lambda \sqrt{-\tilde{g}_{\mu\nu} \frac{dx^\mu}{d\lambda} \frac{dx^\nu}{d\lambda}} \tilde{m}_A$$

i.e. particles follow **geodesics** of $\tilde{g}_{\mu\nu}$ (WEP).

- When self-gravity is not negligible (neutron stars, black holes),

$$S_m = - \sum_A \int d\lambda \sqrt{-\tilde{g}_{\mu\nu} \frac{dx^\mu}{d\lambda} \frac{dx^\nu}{d\lambda}} \tilde{m}_A(\varphi)$$

$\tilde{m}_A(\varphi)$ is a function of the local value of φ to encompass the effect of the background scalar field on the equilibrium of a body. [Eardley 75, DEF 92]

$\tilde{m}_A(\varphi)$ depends on the theory $\mathcal{A}(\varphi)$ and on the EOS of body A.

→ SEP violation

Matter action in the Jordan-frame

$$S_m = - \sum_A \int d\lambda \sqrt{-\tilde{g}_{\mu\nu} \frac{dx^\mu}{d\lambda} \frac{dx^\nu}{d\lambda}} \tilde{m}_A(\varphi)$$

Since $\tilde{g}_{\mu\nu} = \mathcal{A}^2(\varphi)g_{\mu\nu}$:

Matter action in the Einstein-frame

$$S_m = - \sum_A \int d\lambda \sqrt{-g_{\mu\nu} \frac{dx^\mu}{d\lambda} \frac{dx^\nu}{d\lambda}} m_A(\varphi)$$

where we have defined the Einstein-frame mass :

$$m_A(\varphi) \equiv \mathcal{A}(\varphi)\tilde{m}_A(\varphi)$$

The two-body problem hence depends only on 2 fundamental functions, $m_A(\varphi)$ and $m_B(\varphi)$ that encompass **completely** the theory and body-dependence. (GR is recovered when $m_A = cst$, $m_B = cst$.)

I) THE TWO – BODY HAMILTONIAN AT 2PK ORDER

Our starting point : what is known today

Two-body Scalar-Tensor Lagrangian

[DEF 93][Mirshekari, Will 13]

- conservative 2PK dynamics : $\mathcal{O}((\frac{v}{c})^4) \sim \mathcal{O}((\frac{m}{r})^2)$ corrections to Kepler
- Weak field expansion

$$\begin{aligned}g_{\mu\nu} &= \eta_{\mu\nu} + \delta g_{\mu\nu} \\ \varphi &= \varphi_0 + \delta\varphi\end{aligned}$$

- Harmonic coordinates $\partial_\mu(\sqrt{-g}g^{\mu\nu}) = 0$

Generalizes the 2PN GR Lagrangian [Damour, Deruelle 82]

(note that GR dynamics is known at 4PN)

- the fundamental functions $m_A(\varphi)$ and $m_B(\varphi)$ are expanded around φ_0 :

$$\begin{aligned}\ln m_A(\varphi) &\equiv \ln m_A^0 + \alpha_A^0(\varphi - \varphi_0) + \beta_A^0(\varphi - \varphi_0)^2 + \beta_A^{\prime 0}(\varphi - \varphi_0)^3 + \dots \\ \ln m_B(\varphi) &\equiv \ln m_B^0 + \alpha_B^0(\varphi - \varphi_0) + \beta_B^0(\varphi - \varphi_0)^2 + \beta_B^{\prime 0}(\varphi - \varphi_0)^3 + \dots\end{aligned}$$

i.e. the 2PK Lagrangian depends on 8 fundamental **parameters**.

The Jordan-frame Mirshekari-Will Lagrangian is to be translated in terms of the Einstein-frame parametrization introduced above

Two-body 2PK Lagrangian

$$L = -m_A^0 - m_B^0 + L_K + L_{1PK} + L_{2PK} + \dots$$

$$\vec{N} \equiv \frac{\vec{Z}_A - \vec{Z}_B}{R}, \quad \vec{V}_A \equiv \frac{d\vec{Z}_A}{dt}, \quad R \equiv |\vec{Z}_A - \vec{Z}_B|, \quad \vec{A}_A \equiv \frac{d\vec{V}_A}{dt}$$

- Keplerian order :

$$L_K = \frac{1}{2} m_A^0 V_A^2 + \frac{1}{2} m_B^0 V_B^2 + \frac{G_{AB} m_A^0 m_B^0}{R} \quad \text{where} \quad G_{AB} \equiv 1 + \alpha_A^0 \alpha_B^0$$

- post-Keplerian (1PK) :

$$L_{1PK} = \frac{1}{8} m_A^0 V_A^4 + \frac{1}{8} m_B^0 V_B^4 + \frac{G_{AB} m_A^0 m_B^0}{R} \left(\frac{3}{2} (V_A^2 + V_B^2) - \frac{7}{2} \vec{V}_A \cdot \vec{V}_B - \frac{1}{2} (\vec{N} \cdot \vec{V}_A)(\vec{N} \cdot \vec{V}_B) + \bar{\gamma}_{AB} (\vec{V}_A - \vec{V}_B)^2 \right) - \frac{G_{AB}^2 m_A^0 m_B^0}{2R^2} \left(m_A^0 (1 + 2\bar{\beta}_B) + m_B^0 (1 + 2\bar{\beta}_A) \right)$$

$$\text{where} \quad \bar{\gamma}_{AB} \equiv -\frac{2\alpha_A^0 \alpha_B^0}{1 + \alpha_A^0 \alpha_B^0} \quad \bar{\beta}_A \equiv \frac{1}{2} \frac{\beta_A^0 (\alpha_B^0)^2}{(1 + \alpha_A^0 \alpha_B^0)^2} \quad (A \leftrightarrow B)$$

- post-post-Keplerian (2PK) :

$$\begin{aligned}
 L_{2\text{PK}} = & \frac{1}{16} m_A^0 V_A^6 \\
 & + \frac{G_{AB} m_A^0 m_B^0}{R} \left[\frac{1}{8} (7 + 4\bar{\gamma}_{AB}) \left(V_A^4 - V_A^2 (\vec{N} \cdot \vec{V}_B)^2 \right) - (2 + \bar{\gamma}_{AB}) V_A^2 (\vec{V}_A \cdot \vec{V}_B) + \frac{1}{8} (\vec{V}_A \cdot \vec{V}_B)^2 \right. \\
 & \quad \left. + \frac{1}{16} (15 + 8\bar{\gamma}_{AB}) V_A^2 V_B^2 + \frac{3}{16} (\vec{N} \cdot \vec{V}_A)^2 (\vec{N} \cdot \vec{V}_B)^2 + \frac{1}{4} (3 + 2\bar{\gamma}_{AB}) \vec{V}_A \cdot \vec{V}_B (\vec{N} \cdot \vec{V}_A) (\vec{N} \cdot \vec{V}_B) \right] \\
 & + \frac{G_{AB}^2 m_B^0 (m_A^0)^2}{R^2} \left[\frac{1}{8} \left(2 + 12\bar{\gamma}_{AB} + 7\bar{\gamma}_{AB}^2 + 8\bar{\beta}_B - 4\delta_A \right) V_A^2 + \frac{1}{8} \left(14 + 20\bar{\gamma}_{AB} + 7\bar{\gamma}_{AB}^2 + 4\bar{\beta}_B - 4\delta_A \right) V_B^2 \right. \\
 & \quad - \frac{1}{4} \left(7 + 16\bar{\gamma}_{AB} + 7\bar{\gamma}_{AB}^2 + 4\bar{\beta}_B - 4\delta_A \right) \vec{V}_A \cdot \vec{V}_B - \frac{1}{4} \left(14 + 12\bar{\gamma}_{AB} + \bar{\gamma}_{AB}^2 - 8\bar{\beta}_B + 4\delta_A \right) (\vec{V}_A \cdot \vec{N}) (\vec{V}_B \cdot \vec{N}) \\
 & \quad \left. + \frac{1}{8} \left(28 + 20\bar{\gamma}_{AB} + \bar{\gamma}_{AB}^2 - 8\bar{\beta}_B + 4\delta_A \right) (\vec{N} \cdot \vec{V}_A)^2 + \frac{1}{8} \left(4 + 4\bar{\gamma}_{AB} + \bar{\gamma}_{AB}^2 + 4\delta_A \right) (\vec{N} \cdot \vec{V}_B)^2 \right] \\
 & + \frac{G_{AB}^3 (m_A^0)^3 m_B^0}{2R^3} \left[1 + \frac{2}{3} \bar{\gamma}_{AB} + \frac{1}{6} \bar{\gamma}_{AB}^2 + 2\bar{\beta}_B + \frac{2}{3} \delta_A + \frac{1}{3} \epsilon_B \right] + \frac{G_{AB}^3 (m_A^0)^2 (m_B^0)^2}{8R^3} \left[19 + 8\bar{\gamma}_{AB} + 8(\bar{\beta}_A + \bar{\beta}_B) + 4\zeta \right] \\
 & - \frac{1}{8} G_{AB} m_A^0 m_B^0 \left(2(7 + 4\bar{\gamma}_{AB}) \vec{A}_A \cdot \vec{V}_B (\vec{N} \cdot \vec{V}_B) + \vec{N} \cdot \vec{A}_A (\vec{N} \cdot \vec{V}_B)^2 - (7 + 4\bar{\gamma}_{AB}) \vec{N} \cdot \vec{A}_A V_B^2 \right) \\
 & + (A \leftrightarrow B)
 \end{aligned}$$

where $\delta_A \equiv \frac{(\alpha_A^0)^2}{(1 + \alpha_A^0 \alpha_B^0)^2}$ $\epsilon_A \equiv \frac{(\beta'_A \alpha_B^3)^0}{(1 + \alpha_A^0 \alpha_B^0)^3}$ $\zeta \equiv \frac{\beta_A^0 \alpha_A^0 \alpha_B^0 \beta_B^0}{(1 + \alpha_A^0 \alpha_B^0)^3}$ $(A \leftrightarrow B)$

L is written in Harmonic coordinates, and depends on R , \vec{V}_A and on the accelerations \vec{A}_A at 2PK level :

$$L_{2\text{PK}} \ni \boxed{-\frac{1}{8} G_{AB} m_A^0 m_B^0 \left(2(7 + 4\bar{\gamma}_{AB}) \vec{A}_A \cdot \vec{V}_B (\vec{N} \cdot \vec{V}_B) + \vec{N} \cdot \vec{A}_A (\vec{N} \cdot \vec{V}_B)^2 - (7 + 4\bar{\gamma}_{AB}) \vec{N} \cdot \vec{A}_A V_B^2 \right)}$$

→ **Order reduction ?**

contact transformation

[Schäfer 83, Damour-Schäfer 91]

Order reduction : **contact transformation** [Schäfer 83, Damour-Schäfer 91]

1) Add a generic 2PK total time derivative,

$$L \rightarrow L + \frac{df}{dt} \equiv L_f$$

$$\begin{aligned} \frac{f}{m_A^0 m_B^0} \equiv & G_{AB} \left[(f_1 V_A^2 + f_2 \vec{V}_A \cdot \vec{V}_B + f_3 V_B^2)(\vec{N} \cdot \vec{V}_A) - (f_4 V_A^2 + f_5 \vec{V}_A \cdot \vec{V}_B + f_6 V_B^2)(\vec{N} \cdot V_B) \right. \\ & \left. + f_7 (\vec{N} \cdot \vec{V}_A)^3 + f_8 (\vec{N} \cdot \vec{V}_A)^2 (\vec{N} \cdot \vec{V}_B) - f_9 (\vec{N} \cdot \vec{V}_B)^2 (\vec{N} \cdot \vec{V}_A) - f_{10} (\vec{N} \cdot \vec{V}_B)^3 \right] \\ & + G_{AB}^2 \left[f_{11} \left(\frac{m_A^0}{R} \right) (\vec{N} \cdot \vec{V}_A) + f_{12} \left(\frac{m_B^0}{R} \right) (\vec{N} \cdot \vec{V}_A) - f_{13} \left(\frac{m_A^0}{R} \right) (\vec{N} \cdot \vec{V}_B) - f_{14} \left(\frac{m_B^0}{R} \right) (\vec{N} \cdot \vec{V}_B) \right] \end{aligned}$$

where f is a generic function, depending on 14 parameters f_i .

2) Replace the accelerations by their leading on-shell expressions :

$$L_f \rightarrow L_f \left(\vec{A}_A \rightarrow -\vec{N} \frac{G_{AB} m_B^0}{R^2}, \vec{A}_B \rightarrow \vec{N} \frac{G_{AB} m_A^0}{R^2} \right) \equiv L_f^{red}$$

\Leftrightarrow **implicit coordinate change** (contact transformation) : $\vec{Z}_A \rightarrow \vec{Z}_A + \delta \vec{Z}_A$

$$\begin{aligned} \delta \vec{Z}_A = & \frac{G_{AB} m_B^0}{8} \left[2(7 + 4\bar{\gamma}_{AB}) \vec{V}_B (\vec{N} \cdot \vec{V}_B) - \vec{N} \left((7 + 4\bar{\gamma}_{AB}) V_B^2 - (\vec{N} \cdot \vec{V}_B)^2 \right) \right] \\ & - G_{AB} m_B^0 \left[\vec{V}_A \left(2f_1(N \cdot V_A) - 2f_4(N \cdot V_B) \right) + \vec{V}_B \left(f_2(N \cdot V_A) - f_5(N \cdot V_B) \right) \right] \\ & + \vec{N} \left(f_1 V_A^2 + f_2 V_A \cdot V_B + f_3 V_B^2 + 3f_7(N \cdot V_A)^2 + 2f_8(N \cdot V_A)(N \cdot V_B) - f_9(N \cdot V_B)^2 + f_{11} \frac{G_{AB} m_A^0}{R} + f_{12} \frac{G_{AB} m_B^0}{R} \right) \end{aligned}$$

- We have on hand a **whole class of coordinate systems** labeled by 14 parameters f_i for which L_f^{red} is **ordinary**.
- The harmonic coordinates do not belong to this class.

- The two-body Hamiltonians are derived from L_f^{red} through a further **Legendre transformation** :

$$\vec{P}_A = \frac{\partial L_f^{red}}{\partial \vec{V}_A}, \quad \vec{P}_B = \frac{\partial L_f^{red}}{\partial \vec{V}_B}, \quad H = \vec{P}_A \cdot \vec{V}_A + \vec{P}_B \cdot \vec{V}_B - L_f^{red}$$

- In the centre-of-mass frame : $\boxed{\vec{P}_A + \vec{P}_B \equiv \vec{0}}$
i.e. $\vec{Z} \equiv \vec{Z}_A - \vec{Z}_B$ and $\vec{P} \equiv \vec{P}_A = -\vec{P}_B$
- The relative motion is planar \rightarrow use polar coordinates $(Q, P) \equiv (R, \Phi, P_R, P_\Phi)$

General structure of a centre-of-mass frame Hamiltonian $H(Q, P)$

17 coefficients

$$H = M + \left(\frac{P^2}{2\mu} - \mu \frac{G_{AB}M}{R} \right) + H^{1PK} + H^{2PK} + \dots$$

- $\frac{H^{1PK}}{\mu} = \left(h_1^{1PK} \hat{P}^4 + h_2^{1PK} \hat{P}^2 \hat{P}_R^2 + h_3^{1PK} \hat{P}_R^4 \right) + \frac{1}{\hat{R}} \left(h_4^{1PK} \hat{P}^2 + h_5^{1PK} \hat{P}_R^2 \right) + \frac{h_6^{1PK}}{\hat{R}^2}$
- $\frac{H^{2PK}}{\mu} = \left(h_1^{2PK} \hat{P}^6 + h_2^{2PK} \hat{P}^4 \hat{P}_R^2 + h_3^{2PK} \hat{P}^2 \hat{P}_R^4 + h_4^{2PK} \hat{P}_R^6 \right) + \frac{1}{\hat{R}} \left(h_5^{2PK} \hat{P}^4 + h_6^{2PK} \hat{P}_R^2 \hat{P}^2 + h_7^{2PK} \hat{P}_R^4 \right) + \frac{1}{\hat{R}^2} \left(h_8^{2PK} \hat{P}^2 + h_9^{2PK} \hat{P}_R^2 \right) + \frac{h_{10}^{2PK}}{\hat{R}^3}$

$$\mu \equiv \frac{m_A^0 m_B^0}{M}, \quad M \equiv m_A^0 + m_B^0$$

$$10+6+1=17$$

In the Scalar-Tensor case :

The 17 h_i^{MPK} coefficients are computed explicitly and depend on :

- the 14 f_i (coordinate system) parameters
- the 8 fundamental parameters built from $m_A(\varphi)$ and $m_B(\varphi)$

Reminder :

$$\ln m_A(\varphi) \equiv \ln m_A^0 + \alpha_A^0(\varphi - \varphi_0) + \beta_A^0(\varphi - \varphi_0)^2 + \beta_A^{\prime 0}(\varphi - \varphi_0)^3 + \dots$$

$$\ln m_B(\varphi) \equiv \ln m_B^0 + \alpha_B^0(\varphi - \varphi_0) + \beta_B^0(\varphi - \varphi_0)^2 + \beta_B^{\prime 0}(\varphi - \varphi_0)^3 + \dots$$

Recap

- Start from the 2PK two-body Lagrangian (8 parameters)
- Order reduce it through a contact transformation (14 f_i parameters)
- Deduce the 17 h_i^{NPK} coefficients of $H(Q, P)$

$H(Q, P)$ contains all the 2PK information concerning the two-body dynamics at 2PK order, but is **heavy** !

We will contrast it with a much simpler problem, the **geodesic of a test particle in an effective metric**.

II) A TEST PARTICLE IN A SSS EFFECTIVE METRIC

Geodesic motion in a static, spherically symmetric metric

In Schwarzschild-Droste coordinates (equatorial plane $\theta = \pi/2$) :

$$ds_e^2 = -A(r)dt^2 + B(r)dr^2 + r^2 d\phi^2$$

$A(r)$ and $B(r)$ are arbitrary.

- by staticity and spherical symmetry

$$u_t = -A \frac{dt}{d\lambda} \equiv -E, \quad u_\phi = r^2 \frac{d\phi}{d\lambda} \equiv L$$

- 4-velocity normalization

$$u^\mu u_\mu \equiv -1$$

Combining the 3 constants of motion :

Radial EOM ($u \equiv 1/r$)

$$\left(\frac{dr}{d\lambda}\right)^2 = \frac{1}{AB}F(u) \quad \text{with} \quad F(u) \equiv E^2 - A(u)(1 + L^2u^2)$$

- Circular orbits : $F(u) = 0$ and $F'(u) = 0$, hence

$$L^2(u) = -\frac{A'}{(Au^2)'}, \quad E(u) = A\sqrt{\frac{2u}{(Au^2)'}}$$

- Easily yields predictions in the Strong-field regime, e.g. ISCO :

$$F'(u_{\text{ISCO}}) = F''(u_{\text{ISCO}}) = 0 \quad \Rightarrow \quad \boxed{\frac{A''}{A'} = \frac{(Au^2)''}{(Au^2)'}}$$

- One hence easily computes the ISCO orbital frequency

$$\boxed{\omega_{\text{ISCO}} = \frac{d\phi}{dt} = \frac{L}{E} Au^2 \Big|_{u_{\text{ISCO}}}}$$

Note : In these Schwarzschild-Droste coordinates, the dynamics depends only on $A = -g_{00}^e$ for circular orbits

Equivalently, this dynamics is described by the **Lagrangian**

$$L_e = -\mu \sqrt{-g_{\mu\nu}^e \frac{dx^\mu}{dt} \frac{dx^\nu}{dt}} = -\mu \sqrt{A - B\dot{r}^2 - r^2\dot{\phi}^2}$$

Effective Hamiltonian $H_e(q, p)$:

$$H_e(q, p) = \sqrt{A \left(\mu^2 + \frac{p_r^2}{B} + \frac{p_\phi^2}{r^2} \right)} \quad \text{with} \quad p_r \equiv \frac{\partial L_e}{\partial \dot{r}} \quad , \quad p_\phi \equiv \frac{\partial L_e}{\partial \dot{\phi}}$$

Can be **expanded** :

$$\begin{aligned} A(r) &= 1 + \frac{a_1}{r} + \frac{a_2}{r^2} + \frac{a_3}{r^3} + \dots \\ B(r) &= 1 + \frac{b_1}{r} + \frac{b_2}{r^2} + \dots \end{aligned}$$

i.e. depend on **5 effective parameters** at 2PK order, to be determined.

Recap

- On one side, the 2PK two-body Hamiltonian $H(Q, P)$, depending on 17 parameters h_i^{NPK}
- On the other side, a simple effective Hamiltonian $H_e(q, p)$, depending on 5 parameters a_i, b_i .

Can we build a map between both Hamiltonians ?

EOB mapping :

[Buonanno, Damour 98]

Requires imposing a functional relation $H_e = f_{\text{EOB}}(H)$ by means of a canonical transformation

III) THE EOB MAPPING

1) Exploit the power of canonical transformations :

$$H(Q, P) \rightarrow H(q, p)$$

We take as a generic ansatz $G(Q, p)$ that depends on **9 parameters** at 2PK order :

$$G(Q, p) = R p_r \left(\alpha_1 \mathcal{P}^2 + \beta_1 \hat{p}_r^2 + \frac{\gamma_1}{\hat{R}} + \alpha_2 \mathcal{P}^4 + \beta_2 \mathcal{P}^2 \hat{p}_r^2 + \gamma_2 \hat{p}_r^4 + \delta_2 \frac{\mathcal{P}^2}{\hat{R}} + \epsilon_2 \frac{\hat{p}_r^2}{\hat{R}} + \frac{\eta_2}{\hat{R}^2} + \dots \right)$$

$$\mu = \frac{m_A^0 m_B^0}{m_A^0 + m_B^0}, \quad M = m_A^0 + m_B^0$$

$$r(Q, p) = R + \frac{\partial G}{\partial p_r}, \quad \phi(Q, p) = \Phi + \frac{\partial G}{\partial p_\phi}, \quad P_R(Q, p) = p_r + \frac{\partial G}{\partial R}, \quad P_\Phi(Q, p) = p_\phi + \frac{\partial G}{\partial \Phi}$$

- does not depend on time (conservative), nor on Φ (isotropic)
- generates 1PK and higher order coordinate changes

2) Relate H to H_e through a functional relation $H_e = f_{\text{EOB}}(H)$

- at 2PK :

$$\frac{H_e(q, p)}{\mu} - 1 = \left(\frac{H(q, p) - M}{\mu} \right) \left[1 + \frac{\bar{\nu}_1}{2} \left(\frac{H(q, p) - M}{\mu} \right) + \bar{\nu}_2 \left(\frac{H(q, p) - M}{\mu} \right)^2 + \dots \right]$$

the Hamiltonians identifying at Keplerian order.

- In GR, $\bar{\nu}_1 = \nu$ while $\bar{\nu}_2 \dots = 0$ at least up to 4PN

The exact quadratic relation

As proven recently to all orders from PM [Damour 2016]:

$$\frac{H_e(q, p)}{\mu} - 1 = \left(\frac{H(q, p) - M}{\mu} \right) \left[1 + \frac{\nu}{2} \left(\frac{H(q, p) - M}{\mu} \right) \right]$$

where $\nu = \frac{m_A^0 m_B^0}{(m_A^0 + m_B^0)^2}$, $M = m_A^0 + m_B^0$, $\mu = \frac{m_A^0 m_B^0}{M}$

$$\frac{H_e(q, p)}{\mu} - 1 = \left(\frac{H(q, p) - M}{\mu} \right) \left[1 + \frac{\nu}{2} \left(\frac{H(q, p) - M}{\mu} \right) \right]$$

- H_e depends on 5 parameters

$$A(r) = 1 + \frac{a_1}{r} + \frac{a_2}{r^2} + \frac{a_3}{r^3} + \dots, \quad B(r) = 1 + \frac{b_1}{r} + \frac{b_2}{r^2} + \dots$$

- H depends on 17 coefficients (h_i^{NPK});
- The canonical transformation depends on 9 parameters (α_i, β_i, \dots);

$$17 = 9 + 5 + 3$$

Hence, **3** constraints on the h_i^{NPK} coefficients of the two-body Hamiltonian.

→ The two-body problem can be mapped towards a geodesic only for a subclass of theories.

- **At 1PK order, one constraint :**

$$2h_2^{1\text{PK}} + 3h_3^{1\text{PK}} = 0$$

By Lorentz invariance of the kinematical terms $m_A^0 \sqrt{1 - V_A^2}$ at the Lagrangian level , $h_2^{1\text{PK}} = h_3^{1\text{PK}} = 0$ in ST theory.

- **At 2PK order, two constraints :** the first one

$$h_4^{2\text{PK}} = -\frac{2}{45} \left(12h_2^{2\text{PK}} + 18h_3^{2\text{PK}} + (h_2^{1\text{PK}})^2 \right)$$

is no more restrictive.

However, the second one

$$\begin{aligned} h_1^{2\text{PK}} + \frac{7}{3}h_2^{2\text{PK}} + h_3^{2\text{PK}} + h_5^{2\text{PK}} + h_6^{2\text{PK}} + h_7^{2\text{PK}} = \\ -\frac{h^K}{128}(5 + 2\nu + 5\nu^2) + \frac{1}{8}(1 + \nu)\left((3h_1^{1\text{PK}} + h_2^{1\text{PK}})h^K + h_4^{1\text{PK}} + h_5^{1\text{PK}}\right) + \frac{5}{2}h_1^{1\text{PK}}\left(7h_1^{1\text{PK}}h^K + 2(h_4^{1\text{PK}} + h_5^{1\text{PK}})\right) \\ + \frac{1}{6}h_2^{1\text{PK}}\left(13h_2^{1\text{PK}}h^K + 10(h_4^{1\text{PK}} + h_5^{1\text{PK}})\right) + \frac{35}{3}h_1^{1\text{PK}}h_2^{1\text{PK}}h^K, \end{aligned}$$

is restrictive.

- satisfied by the Scalar-Tensor coefficients, for any $f_i \dots$
- but not by Electrodynamics.

In **Scalar-Tensor theories**, injecting the expressions of h_i^{MPK} , the identification

$$\frac{H_e(q, p)}{\mu} - 1 = \left(\frac{H(q, p) - M}{\mu} \right) \left[1 + \frac{\nu}{2} \left(\frac{H(q, p) - M}{\mu} \right) \right]$$

- yields a **unique** solution for H_e that does not depend on the f_i parameters (covariance).
- H_e contain all the 2PK physical information contained in $H(Q, P)$.

Part of the **complexity** of $H(Q, P)$, e.g. f , is **hidden in the canonical transformation** :

$$G(Q, p) = R p_r \left(\alpha_1 \mathcal{P}^2 + \beta_1 \hat{p}_r^2 + \frac{\gamma_1}{\hat{R}} + \alpha_2 \mathcal{P}^4 + \beta_2 \mathcal{P}^2 \hat{p}_r^2 + \gamma_2 \hat{p}_r^4 + \delta_2 \frac{\mathcal{P}^2}{\hat{R}} + \epsilon_2 \frac{\hat{p}_r^2}{\hat{R}} + \frac{\eta_2}{\hat{R}^2} + \dots \right)$$

whose **9** parameters read :

$$\begin{aligned} \alpha_1 &= -\frac{\nu}{2}, \quad \beta_1 = 0, \quad \gamma_1 = G_{AB} \left(1 + \bar{\gamma}_{AB} + \frac{\nu}{2} \right), \quad \alpha_2 = \frac{1}{8}(1 - \nu)\nu, \quad \beta_2 = 0, \quad \gamma_2 = \frac{\nu^2}{2}, \\ \delta_2 &= G_{AB} \left[f_6 \frac{m_A^0}{M} + f_1 \frac{m_B^0}{M} - \nu \left(f_1 + f_6 + (-f_3 + f_5 + f_6) \frac{m_A^0}{M} + (f_1 + f_2 - f_4) \frac{m_B^0}{M} - \frac{3}{2}(1 + \bar{\gamma}_{AB}) + \frac{\nu}{8} \right) \right], \\ \epsilon_2 &= G_{AB} \left[-\frac{\nu^2}{8} + f_{10} \frac{m_A^0}{M} + f_7 \frac{m_B^0}{M} - \nu \left(f_7 + f_{10} + (f_9 + f_{10}) \frac{m_A^0}{M} + (f_7 + f_8) \frac{m_B^0}{M} \right) \right], \\ \eta_2 &= \frac{1}{8} G_{AB}^2 \left[8\langle \bar{\beta} \rangle - 4\langle \delta \rangle + 4\bar{\gamma}_{AB} + 3\bar{\gamma}_{AB}^2 + \nu \left(-38 + 4(\bar{\beta}_A + \bar{\beta}_B) - 24\bar{\gamma}_{AB} + 2\nu \right) \right] \\ &\quad + G_{AB}^2 \left(f_{13} \frac{m_A^0}{M} + f_{12} \frac{m_B^0}{M} + \nu(f_{11} - f_{12} - f_{13} + f_{14}) \right) \end{aligned}$$

$$ds_e^2 = -A(r)dt + B(r)dr^2 + r^2 d\theta^2$$

Scalar-Tensor effective metric

$$A(r) = 1 - 2 \left(\frac{G_{AB} M}{r} \right) + 2 \left[\langle \bar{\beta} \rangle - \bar{\gamma}_{AB} \right] \left(\frac{G_{AB} M}{r} \right)^2 + \left[2\nu + \delta a_3^{\text{ST}} \right] \left(\frac{G_{AB} M}{r} \right)^3 + \dots$$

$$B(r) = 1 + 2 \left[1 + \bar{\gamma}_{AB} \right] \left(\frac{G_{AB} M}{r} \right) + \left[2(2 - 3\nu) + \delta b_2^{\text{ST}} \right] \left(\frac{G_{AB} M}{r} \right)^2 + \dots$$

Consistent with General Relativity : $m_A(\varphi) = cst$ yields back

General Relativity 2PN effective metric

[Buonanno, Damour 98]

$$A_{\text{GR}}(r) = 1 - 2 \left(\frac{G_* M}{r} \right) + 2\nu \left(\frac{G_* M}{r} \right)^3 + \dots$$

$$B_{\text{GR}}(r) = 1 + 2 \left(\frac{G_* M}{r} \right) + 2(2 - 3\nu) \left(\frac{G_* M}{r} \right)^2 + \dots$$

(i) The “bare” gravitational constant G_* is replaced by the effective one

$$G_* \rightarrow G_{AB} \equiv 1 + \alpha_A^0 \alpha_B^0$$

at all orders.

(ii) At 1PK level,

$$A(r) = 1 - 2 \left(\frac{G_{AB} M}{r} \right) + 2 \left[\langle \bar{\beta} \rangle - \bar{\gamma}_{AB} \right] \left(\frac{G_{AB} M}{r} \right)^2 + \dots$$

$$B(r) = 1 + 2 \left[1 + \bar{\gamma}_{AB} \right] \left(\frac{G_{AB} M}{r} \right) + \dots$$

one recognizes the **PPN Eddington metric** written in Droste coordinates, with :

$$\beta^{\text{Edd}} = 1 + \langle \bar{\beta} \rangle, \quad \gamma^{\text{Edd}} = 1 + \bar{\gamma}_{AB}$$

Where

$$\langle \bar{\beta} \rangle \equiv \frac{m_A^0 \bar{\beta}_B + m_B^0 \bar{\beta}_A}{m_A^0 + m_B^0} \quad \bar{\gamma}_{AB} \equiv -\frac{2\alpha_A^0 \alpha_B^0}{1 + \alpha_A^0 \alpha_B^0} \quad \bar{\beta}_A \equiv \frac{1}{2} \frac{\beta_A^0 (\alpha_B^0)^2}{(1 + \alpha_A^0 \alpha_B^0)^2}$$

Scalar-Tensor effective metric

$$A(r) = 1 - 2 \left(\frac{G_{AB} M}{r} \right) + 2 \left[\langle \bar{\beta} \rangle - \bar{\gamma}_{AB} \right] \left(\frac{G_{AB} M}{r} \right)^2 + \left[2\nu + \delta a_3^{\text{ST}} \right] \left(\frac{G_{AB} M}{r} \right)^3 + \dots$$

$$B(r) = 1 + 2 \left[1 + \bar{\gamma}_{AB} \right] \left(\frac{G_{AB} M}{r} \right) + \left[2(2 - 3\nu) + \delta b_2^{\text{ST}} \right] \left(\frac{G_{AB} M}{r} \right)^2 + \dots$$

(iii) 2PK corrections

$$\delta a_3^{\text{ST}} \equiv \frac{1}{12} \left[-20\bar{\gamma}_{AB} - 35\bar{\gamma}_{AB}^2 - 24\langle \bar{\beta} \rangle (1 - 2\bar{\gamma}_{AB}) + 4(\langle \delta \rangle - \langle \epsilon \rangle) \right. \\ \left. + \nu \left(-36(\bar{\beta}_A + \bar{\beta}_B) + 4\bar{\gamma}_{AB}(10 + \bar{\gamma}_{AB}) + 4(\epsilon_A + \epsilon_B) + 8(\delta_A + \delta_B) - 24\zeta \right) \right]$$

$$\delta b_2^{\text{ST}} \equiv \left[4\langle \bar{\beta} \rangle - \langle \delta \rangle + \bar{\gamma}_{AB} \left(9 + \frac{19}{4} \bar{\gamma}_{AB} \right) + \nu \left(2\langle \bar{\beta} \rangle - 4\bar{\gamma}_{AB} \right) \right]$$

$$\delta_A \equiv \frac{(\alpha_A^0)^2}{(1 + \alpha_A^0 \alpha_B^0)^2} \quad \epsilon_A \equiv \frac{(\beta'_A \alpha_B^3)^0}{(1 + \alpha_A^0 \alpha_B^0)^3} \quad \zeta \equiv \frac{\beta_A^0 \alpha_A^0 \alpha_B^0 \beta_B^0}{(1 + \alpha_A^0 \alpha_B^0)^3}$$

Recap :

- By means of a canonical transformation and a quadratic relation

$$\frac{H_e(q, p)}{\mu} - 1 = \left(\frac{H(q, p) - M}{\mu} \right) \left[1 + \frac{\nu}{2} \left(\frac{H(q, p) - M}{\mu} \right) \right]$$

determined uniquely the effective Hamiltonian H_e .

- The whole 2PK dynamics has been reduced to the simple geodesic of an effective metric :

$$A(r) = 1 - 2 \left(\frac{G_{AB} M}{r} \right) + 2 \left[\langle \bar{\beta} \rangle - \bar{\gamma}_{AB} \right] \left(\frac{G_{AB} M}{r} \right)^2 + \left[2\nu + \delta a_3^{\text{ST}} \right] \left(\frac{G_{AB} M}{r} \right)^3 + \dots$$

$$B(r) = 1 + 2 \left[1 + \bar{\gamma}_{AB} \right] \left(\frac{G_{AB} M}{r} \right) + \left[2(2 - 3\nu) + \delta b_2^{\text{ST}} \right] \left(\frac{G_{AB} M}{r} \right)^2 + \dots$$

IV) ST – PARAMETRISED EOB DYNAMICS

- The inversion of the $H_e = f_{\text{EOB}}(H)$ hence defines a unique, “resummed” EOB Hamiltonian :

$$H_{\text{EOB}} = M \sqrt{1 + 2\nu \left(\frac{H_e}{\mu} - 1 \right)}$$

The dynamics deduced from H_{EOB} and the “real” Hamiltonians H are, by construction, equivalent up to 2PK order.

- H_{EOB} hence defines a resummed dynamics, that may capture some features of the strong field regime.

$$H_{\text{EOB}} = M \sqrt{1 + 2\nu \left(\frac{H_e}{\mu} - 1 \right)}, \quad \text{where} \quad H_e = \sqrt{A \left(\mu^2 + \frac{p_r^2}{B} + \frac{p_\phi^2}{r^2} \right)}$$

But H_{EOB} and H_e are conservative :

$$\Rightarrow \left(\frac{\partial H_{\text{EOB}}}{\partial H_e} \right) = \frac{1}{\sqrt{1 + 2\nu(E - 1)}} \quad \text{since} \quad H_e = \mu E \quad \text{on-shell}$$

Hence the two-body eom, deduced from H_{EOB} are identical to the effective ones, deduced from H_e , to within a simple time rescaling

$$t \rightarrow t \sqrt{1 + 2\nu(E - 1)}$$

In particular, the **orbital frequency**, deduced from H_{EOB} , is

$$\Omega = \frac{\partial H_{\text{EOB}}}{\partial H_e} \frac{\partial H_e}{\partial p_\phi} = \frac{j u^2 A}{G_{AB} M E \sqrt{1 + 2\nu(E - 1)}} \quad u = \frac{G_{AB} M}{r}$$

where, for circular orbits

$$j^2(u) = -\frac{A'}{(Au^2)'} , \quad E(u) = A \sqrt{\frac{2u}{(Au^2)'}}$$

and can be in particular evaluated **at the level of the ISCO**, u_{ISCO} , such that :

$$\frac{A''}{A'} = \frac{(Au^2)''}{(Au^2)'}$$

→ **ST corrections to the ISCO location and frequency ?**
(typical near merger)

The ISCO location and frequency depend only on $A(u) = -g_{00}^e$

Last ingredient : the ST-corrected $A(u; \nu)$

$$u \equiv \frac{G_{AB} M}{r}, \quad \nu = \frac{m_A^0 m_B^0}{(m_A^0 + m_B^0)^2}$$

ST-corrected $A(u; \nu)$

$$A(u; \nu) = A_{2\text{PN}}^{\text{GR}}(u; \nu) + 2\epsilon_{1\text{PK}} u^2 + (\epsilon_{2\text{PK}}^0 + \nu \epsilon_{2\text{PK}}^\nu) u^3$$

where

$$\epsilon_{1\text{PK}} \equiv \langle \bar{\beta} \rangle - \bar{\gamma}_{AB}$$

$$\epsilon_{2\text{PK}}^0 \equiv \frac{1}{12} \left[-20\bar{\gamma}_{AB} - 35\bar{\gamma}_{AB}^2 - 24\langle \bar{\beta} \rangle (1 - 2\bar{\gamma}_{AB}) + 4(\langle \delta \rangle - \langle \epsilon \rangle) \right]$$

$$\epsilon_{2\text{PK}}^\nu \equiv -3(\bar{\beta}_A + \bar{\beta}_B) + \frac{1}{3}\bar{\gamma}_{AB}(10 + \bar{\gamma}_{AB}) + \frac{1}{3}(\epsilon_A + \epsilon_B) + \frac{2}{3}(\delta_A + \delta_B) - 2\zeta$$

ST-Corrections described by 3 parameters, $(\epsilon_{1\text{PK}}, \epsilon_{2\text{PK}}^0, \epsilon_{2\text{PK}}^\nu)$

- **BUT** numerically driven by $(\alpha_A^0)^2$ (c.f. DEF, diagrammatic methods)
- When $(\alpha_A^0)^2 \ll 1$, $\epsilon_{1\text{PK}} \sim \epsilon_{2\text{PK}}^0 \sim \epsilon_{2\text{PK}}^\nu$ and ST-corrections are perturbative

In this perturbative approach, **best available EOB-NR function** for GR :

$$A_{2\text{PN}}^{\text{GR}}(u; \nu) \rightarrow \boxed{A_{\text{EOBNR}}^{\text{GR}}(u; \nu) = \mathcal{P}_5^1[A_{5\text{PN}}^{\text{Taylor}}]}$$

i.e. the (1, 5) Padé approximant of the truncated 5PN expansion :

$$A_{5\text{PN}}^{\text{Taylor}} = 1 - 2u + 2\nu u^3 + \nu a_4 u^4 + (a_5^c + a_5^{\text{ln}} \ln u) u^5 + \nu (a_6^c + a_6^{\text{ln}} \ln u) u^6$$


[Damour, Nagar, Reisswig, Pollney 2016]

- smoothly connected to Schwarzschild when $\nu \rightarrow 0$
- $a_6^c(\nu)$ is obtained by calibration with Numerical Relativity


ISCO Locations and frequency, equal-mass case ($\nu = 1/4$)

- **1PK corrections**, $A = A_{\text{EOBNR}}^{\text{GR}}(u; \nu) + 2\epsilon_{1\text{PK}}u^2$
 - **2PK corrections**, $A = A_{\text{EOBNR}}^{\text{GR}}(u; \nu) + 2\epsilon_{1\text{PK}}u^2 + (\epsilon_{2\text{PK}}^0 + \nu\epsilon_{2\text{PK}}^\nu)u^3$
- setting $\epsilon_{2\text{PK}}^0 + \nu\epsilon_{2\text{PK}}^\nu \equiv \epsilon_{1\text{PK}}$

$u_{\text{ISCO}}(\epsilon_{1\text{PK}})$




$G_{\text{AB}}M\Omega_{\text{ISCO}}(\epsilon_{1\text{PK}})$



→ dramatic increase when $\epsilon_{1\text{PK}} \sim 10^{-1}$

$$A(u) \quad (\nu = 1/4, \epsilon_{1\text{PK}} = 0.08)$$



- **2PK corrections**, $A = A_{\text{EOBNR}}^{\text{GR}}(u; \nu) + 2\epsilon_{1\text{PK}}u^2 + (\epsilon_{2\text{PK}}^0 + \nu\epsilon_{2\text{PK}}^\nu)u^3$

- **2PK Padeed corrections**,



$$A = \mathcal{P}_5^1[A_{\text{EOBNR}}^{\text{GR}}(u; \nu) + 2\epsilon_{1\text{PK}}u^2 + (\epsilon_{2\text{PK}}^0 + \nu\epsilon_{2\text{PK}}^\nu)u^3]$$

Tiny corrections to A , but significant corrections to the ISCO frequency

ISCO Locations and frequency, equal-mass case ($\nu = 1/4$)

- 2PK Padeed corrections,

$$A = \mathcal{P}_5^1[A_{\text{EOBNR}}^{\text{GR}}(u; \nu) + 2\epsilon_{1\text{PK}}u^2 + (\epsilon_{2\text{PK}}^0 + \nu\epsilon_{2\text{PK}}^\nu)u^3]$$



$$\left. \frac{d(G_{AB}M\Omega)_{\text{ISCO}}}{d\epsilon_{1\text{PK}}} \right|_{\nu=1/4} \simeq 0.13$$

relative correction to GR significant ($\sim 10\%$) when $\epsilon_{1\text{PK}} \sim 10^{-2} - 10^{-1}$

Concluding remarks :

- Remarkably, the EOB approach is valid beyond the scope of General Relativity. In **Scalar-Tensor theories** :

$$A^{2\text{PK}}(u) \equiv \mathcal{P}_5^1[A_{5\text{PN}}^{\text{Taylor}} + 2\epsilon_{1\text{PK}}u^2 + (\epsilon_{2\text{PK}}^0 + \nu\epsilon_{2\text{PK}}^\nu)u^3]$$

- But also applicable for **any theory** whose coefficients h_i^{MPK} satisfy the 3 mapping conditions.
- The Scalar-Tensor example suggests a generic 2PK ansatz

$$A^{\text{PEOB}}(u) \equiv \mathcal{P}_5^1[A_{5\text{PN}}^{\text{Taylor}} + 2(\epsilon_{1\text{PK}}^0 + \nu\epsilon_{1\text{PK}}^\nu)u^2 + (\epsilon_{2\text{PK}}^0 + \nu\epsilon_{2\text{PK}}^\nu)u^3]$$

where $\epsilon_{1\text{PK}}^0$, $\epsilon_{1\text{PK}}^\nu$, $\epsilon_{2\text{PK}}^0$, and $\epsilon_{2\text{PK}}^\nu$ are theory-agnostic Parametrized EOB (PEOB) coefficients.

- The EOB approach has been extended to neutron stars including **tidal effects** (TEOB, [Damour, Nagar 2010]) through 5PN $\mathcal{O}(u^6)$ corrections to $A(u)$. To be compared with ST-EOB $\mathcal{O}(u^2)$.
- Binary pulsar experiments have put **stringent constraints on ST theories** (no dipolar radiation)

$$(\alpha_A^0)^2 < 4 \times 10^{-6}$$

For **any** body A, regardless of its EOS or self-gravity.

- The ISCO ST-correction (significant for $(\alpha_A^0)^2 \gtrsim 10^{-2}$) seems unlikely to improve binary pulsar constraints.

However :

- However, stars subject to dynamical scalarization can develop non perturbative $(\alpha_A^0)^2$ near merger [Barausse, Palenzuela, Ponce, Lehner 2013]. EOB well-suited to investigate this regime !
- The interferometers LIGO-Virgo or even LISA are designed to detect highly redshifted sources. Cosmological history of ST theories ?

Black holes :

- Are known in these ST theories to carry no scalar hair : $m_A(\varphi) = cst$ i.e. no deviation to GR.
- Induce scalar hair by means of $V(\varphi)$ or (vector) gauge field ? (at least 1PK by Lorentz invariance).
- Conditions of no hair theorems may not apply anymore in the strong field, highly dynamical regime of a merger, which is **precisely** explored by the EOB approach !