

PAPER • OPEN ACCESS

Standard model as the topological material

To cite this article: G E Volovik and M A Zubkov 2017 *New J. Phys.* **19** 015009

View the [article online](#) for updates and enhancements.

Related content

- [From standard model of particle physics to room-temperature superconductivity](#)
G E Volovik
- [Anomalous Hall effect in Weyl superconductors](#)
G Bednik, A A Zyuzin and A A Burkov
- [A facility to search for hidden particles at the CERN SPS: the SHiP physics case](#)
Sergey Alekhin, Wolfgang Altmannshofer, Takehiko Asaka et al.

Recent citations

- [Extracting topological information from momentum space propagators](#)
Fabrizio Canfora *et al*
- [Dai-Freed anomalies in particle physics](#)
Iñaki García-Etxebarria and Miguel Montero
- [Chiral Torsional Effect](#)
Z. V. Khaidukov and M. A. Zubkov



PAPER

Standard model as the topological material

OPEN ACCESS

RECEIVED

23 August 2016

REVISED

11 December 2016

ACCEPTED FOR PUBLICATION

6 January 2017

PUBLISHED

23 January 2017

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](#).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

G E Volovik^{1,2,6} and M A Zubkov^{3,4,5}¹ Low Temperature Laboratory, School of Science and Technology, Aalto University, Finland² L.D. Landau Institute for Theoretical Physics, Moscow, Russia³ LE STUDIUM, Loire Valley Institute for Advanced Studies, Tours and Orleans, F-45000 Orleans France⁴ Laboratoire de Mathematiques et de Physique Theorique, Universite de Tours, F-37200 Tours, France⁵ Institute for Theoretical and Experimental Physics, B. Chermushkinskaya 25, Moscow, 117259, Russia⁶ Author to whom any correspondence should be addressed.E-mail: volovik@ltd.tkk.fi

Keywords: Weyl fermions, Dirac fermions, topological stability, standard model

Abstract

The study of the Weyl and Dirac topological materials (topological semimetals, insulators, superfluids and superconductors) opens the route for the investigation of the topological quantum vacua of relativistic fields. The symmetric phase of the standard model (SM), where both electroweak and chiral symmetry are not broken, represents the topological semimetal. The vacua of the SM (and its extensions) in the phases with broken electroweak symmetry represent the topological insulators of different types. We discuss in detail the topological invariants in both the symmetric and broken phases and establish their relation to the stability of vacuum.

1. Introduction

The massless (gapless) Weyl fermions in the symmetric phase of the standard model (SM) of fundamental interactions have common topological properties with the Weyl and Dirac fermions in topological semimetals. The topological stability of the Weyl node in the spectrum of neutrino was first considered in [1], see also [2]. Later the topological invariant for the Weyl points was expressed in terms of the fermionic Green's function [3], and then the topological approach was extended by Hořava to the other types of nodes in the fermionic spectrum, such as Dirac nodal lines and Fermi surfaces [4]; this topological classification of the possible types of zeroes in the spectrum was based on the K -theory. Topological classification then has been extended to the other topological phases of matter—the fully gapped states, such as topological insulators, topological superconductors and the phase B of superfluid ^3He , see [5–8].

The systems (vacua) with the Weyl points both in condensed matter and in particle physics have many exotic properties, such as chiral anomaly. For example, the Adler–Bell–Jackiw equation, which describes the anomalous production of fermions from vacuum [9–11] has been verified in experiments with skyrmions in the chiral superfluid $^3\text{He-A}$ [12], see also [13]. Weyl fermions in semimetals have been considered by Abrikosov and Beneslavskii in 1970 [14]; for the recent reviews on Weyl fermions in semimetals, superconductors and superfluids see [15–19].

In the topological classification an important role is played by the symmetry of the vacuum. This also concerns the symmetric phase of the SM, where both electroweak and chiral symmetries vacuum are not broken. In this phase due to the equal number of the left-handed and the right-handed particles (if the sterile neutrino is included), and due to Lorentz invariance, the total topological charge in the Fermi point situated at $\mathbf{p} = 0$ is zero. Therefore, the topology of the Weyl fermions in the SM is to be supported by symmetry⁷. The modification of the momentum space topological invariants associated with various elements of the SM gauge group has been suggested in [13].

⁷ It is possible that the Weyl points of the SM originate from the nontrivial topology of the underlying vacuum with Majorana fermions [20–22].

In this paper we consider the complete set of the topological invariants for the Fermi point of the SM fermions. We demonstrate that the generating functional for those invariants possesses the Z_6 symmetry, which, relates the elements of the gauge group giving rise to equivalent topological invariants. This is the same Z_6 symmetry of the fermionic representations of the SM, which was discussed in [23]. The existence of this Z_6 symmetry explains, in particular, why the topological invariant that protects all massless SM fermions may be expressed through either the hypercharge generator or through one of the generators of the $SU(2)_L$ subgroup of the SM gauge group (see section 12.3 of [13]).

Next, we discuss the vacua of the SM (and its extensions) in the phases with broken electroweak symmetry. These phases represent the topological insulators of different types. First we consider the conditions at which the parity breaking interactions may be neglected. This in particular requires that the temperature be much smaller than the masses of the corresponding fermions. Under this limit the topological classes of the SM vacua are classified according to the topological invariant associated with the matrix of CT symmetry (the combination of Charge conjugation and Time reversal symmetries), which protects the number of massive Dirac fermions. Notice that if the interactions are neglected at all, the vacuum of the SM in the massive phases would be described by the same topology as the fully gapped superfluid $^3\text{He-B}$ [24]. In superfluid $^3\text{He-B}$ the topological invariant is protected by the chiral symmetry of the system. The CT symmetry in the SM plays a similar role. Presumably, the corresponding topological invariant is relevant for the topological classification of the vacua of the SM at low enough temperatures (pressure, chemical potential etc).

In addition to the topological invariant protected by CT symmetry there exists the topological invariant protected by T-symmetry, which is relevant for the consideration of the SM, when the interactions that break CP (the combination of Charge conjugation and Parity symmetries) are taken into account. This invariant becomes important when the emphasis is on the consideration of the Higgs sector of the SM. The topologically nontrivial phase appears, when the Majorana masses of the left-handed neutrinos are present, the number of which is protected by this topological invariant. We demonstrate that in the noninteracting case of the massive SM Dirac fermions the value of the symmetry protected topological invariant associated with T is equal to zero, $N_{K_T} = 0$. At the same time in the extensions of the SM with the type II neutrino seesaw the value of the topological invariant N_{K_T} (supported by the T-symmetry) is nonzero. Therefore, the phases with and without Majorana masses of the left-handed neutrinos cannot be continuously connected and are, indeed, the different phases. However, we obtain that the type I seesaw is topologically trivial and its vacuum may be transformed without the phase transition to the conventional vacuum of the SM with Dirac neutrino masses.

Depending on the external conditions (temperature, pressure, chemical potentials of various types, etc) the SM and its extensions may exist in various phases. For example, in addition to the ordinary baryonic phase, which is realized at the vanishing temperature, pressure and baryonic chemical potential, in QCD there exist various other phases: several color superconducting phases, the quark–gluon plasma phase, etc [52–55]. The Weinberg–Salam model is typically considered in the two phases: the symmetric high temperature phase with the restored chiral symmetry and the broken low temperature phase with the spontaneously broken $SU(2) \otimes U(1)$ symmetry. The complete SM (containing Quantum Chromodynamics and the Weinberg–Salam model) may possess new phases, which have not been considered yet, at certain external conditions. Various extensions of the SM like the models with Majorana masses of neutrinos, models with several Higgs bosons, models with composite Higgs bosons may also exist in several exotic phases, which have not been considered so far. The momentum space topological invariants discussed in this paper may be applied to the consideration of the phase transitions between the phases of the SM (and its extensions) mentioned above. For the previous consideration of the topologically nontrivial vacua in relativistic quantum field theories based on the topological invariants in momentum space see, for example, [4, 13, 42–48, 50, 51].

The paper is organized as follows. In section 2 we consider the symmetric phase of the SM with unbroken chiral and electroweak symmetries as the phase of the topological semimetal. The complete set of the topological invariants protecting the Weyl points is defined, and the Z_6 symmetry of the corresponding generating functional is established. In sections 3 and 4 we discuss the SM at low temperatures, which are smaller than the mass of the lightest fermion. In section 3 we discuss the situation at the sufficiently small values of pressure and chemical potentials, so that the parity breaking interactions are to be neglected in the consideration of the questions of the stability of vacuum. In section 4 we discuss the topological invariant of the SM and its extensions that remain at work if the parity breaking interactions are taken into account while the CP breaking is neglected. In particular, it is demonstrated that the vacuum with Dirac fermions is topologically trivial (with respect to the invariant protected by time reversal symmetry). At the same time, the vacuum with Majorana masses may be topologically nontrivial.

2. SM in the symmetric phase as the topological semimetal

2.1. Topological invariant for massless fermions

In its gapless (massless) phase the SM belongs to the class of the 3+1 dimensional vacua, which are characterized by the Weyl points in momentum space. The Weyl point is characterized by the momentum space topological invariant N_3 , which protects the masslessness of the fermionic spectrum. The topological invariant for the isolated Weyl point is expressed as the integral of the three-form in terms of the two-point Green's function \mathcal{G} determined in the 4D momentum-frequency space [3, 13]:

$$N_3 = \text{tr } \mathcal{N}, \quad \mathcal{N} = \frac{1}{24\pi^2} \epsilon_{\mu\nu\lambda\gamma} \int_{\sigma} dS^{\gamma} \mathcal{G} \partial_{p_{\mu}} \mathcal{G}^{-1} \mathcal{G} \partial_{p_{\nu}} \mathcal{G}^{-1} \mathcal{G} \partial_{p_{\lambda}} \mathcal{G}^{-1}. \quad (1)$$

The Green's function is an $n \times n$ matrix. For a single species of Weyl fermions one has $n = 2$, and the Green's function is expressed in terms of the Pauli matrices. For general topological condensed matter system the $n \times n$ matrix contains Pauli matrices for spin and for the Bogoliubov spin, and also the crystal band indices of fermions. In particle physics, the $n \times n$ matrix includes Weyl or Dirac matrices and indices of different fermionic species (quarks and leptons of different generations). In SM with 16 species in one generation has $n = 32g$, where g is the number of generations. If expressed in terms of Majorana fermions, the matrix has $n = 64g$. The definition of the Green's function in terms of the functional integral over the fields is given in appendix A. The integral in equation (1) is over the S^3 surface σ embracing the point in the 4D space $\mathbf{p} = 0$, $p_4 = 0$, where p_4 is the frequency along the imaginary axis; tr is the trace over the fermionic indices. For a single species of right-handed Weyl fermions one has $N_3 = 1$, and $N_3 = -1$ for the left-handed Weyl fermions.

It is worth mentioning that the symmetric phase of the SM appears at finite temperatures, while the topological classification and topological invariants are formally applicable only to the ground state (vacuum) of the system. Actually the consideration is valid if the temperature T is much smaller than the characteristic high energy scale of the system $T \ll T_{\text{uv}}$. Here T_{uv} is the scale, at which the SM of fundamental interactions already does not work, and new fields and interactions appear. In this limit all the properties of the systems related to topology, such as chiral anomaly, are determined by these topological invariants. Thus in spite of the fact that the topological invariant N_3 is defined typically for the zero temperature, in the SM this invariant appears to be well-defined at the temperatures above the electroweak transition if those temperatures are smaller than the scale of the ultraviolet completion of the SM, which is at least one order of magnitude higher than the electroweak scale ~ 100 GeV.

If sterile right-handed neutrinos are present in the SM, the number of the left- and the right-handed fermions is equal, $n_{\text{left}} = n_{\text{right}} = 8g$, where g is the number of generations. This is required, for example, if we assume that the lattice regularization is used, where the numbers of the left-handed and the right-handed fermions are equal due to the Nielsen–Ninomiya theorem. Then the trace in equation (1) over all the fermionic species gives the trivial value for the topological invariant, $N_3 = n_{\text{right}} - n_{\text{left}} = 0$. Nevertheless, the vacuum of the SM is topologically nontrivial, because its topology is supported by the symmetry of the SM in the symmetric phase. The $SU(3) \otimes SU(2) \otimes U(1)$ symmetry allows one to introduce the generating function of topological invariants, which contains the powers of the hypercharge Y , the generators of $SU(2)_L$ and $SU(3)_c$:

$$N(\theta_Y, \theta_W^a, \theta_c^i) = \text{tr} [e^{i\theta_W^a \mathcal{W}_a} e^{i\theta_Y \mathcal{Y}} e^{i\theta_c^i C_i} \mathcal{N}] \quad (2)$$

(Here \mathcal{W}_a , $a = 1, 2, 3$ are the generators of $SU(2)_L$ while C_i , $i = 1, \dots, 8$ are the generators of $SU(3)_c$.)

2.2. Z_6 symmetry of the fermionic representations in the SM

Notice that equation (2) obeys the Z_6 invariance (see also [25]):

$$\theta_Y \rightarrow \theta_Y + 2\pi N, \quad e^{i\theta_W^a \mathcal{W}_a} \rightarrow e^{i\theta_W^a \mathcal{W}_a} \times e^{i\pi N}, \quad e^{i\theta_c^i C_i} \rightarrow e^{i\theta_c^i C_i} \times e^{\frac{2\pi i}{3} N} \quad (3)$$

where N is the integer. This invariance might actually mean that the gauge group of the SM is $SU(3) \otimes SU(2) \otimes U(1)/Z_6$ rather than $SU(3) \otimes SU(2) \otimes U(1)$ [23]. The given Z_6 symmetry follows from the assignment of the hypercharges, weak charges and electric charges $Q = Y + W$ of the fermions given by

Fermion	W	Y	Q
$u_L(3)$	$+\frac{1}{2}$	$\frac{1}{6}$	$\frac{2}{3}$
$u_R(3)$	0	$\frac{2}{3}$	$\frac{2}{3}$
$d_L(3)$	$-\frac{1}{2}$	$\frac{1}{6}$	$-\frac{1}{3}$
$d_R(3)$	0	$-\frac{1}{3}$	$-\frac{1}{3}$
ν_L	$+\frac{1}{2}$	$-\frac{1}{2}$	0
ν_R	0	0	0
e_L	$-\frac{1}{2}$	$-\frac{1}{2}$	-1
e_R	0	-1	-1

(4)

According to this table in the SM including strong interactions the group $U(1) \times SU(2) \times SU(3)$ has the global $Z_6 = Z_2 \times Z_3$ -subgroup of elements which act on the SM fermions as an identity element (see equations (61)–(64) in [2] and [23, 25]). This group consists of the following elements g^k :

$$g^k = [e^{i2\pi C_8} e^{2\pi i Y} e^{2\pi i W_3}]^k, \quad k = 1, \dots, 6. \quad (5)$$

where C_8 will be specified below in equation (9). Notice that the Z_6 symmetry of the fermionic representations of the SM takes place in any phases, not only in the symmetric phase. Its elements of equation (5) being applied to any fermion of the SM give 1. In the other words, all SM fermions represent the eigenvectors of the elements of Z_6 corresponding to the eigenvalues equal to unity.

2.3. Maximal number of Dirac massless fermions protected by the topological invariants

The generators \mathcal{W}_a , C_i , \mathcal{Y} commute with the fermion Green's functions taken in the Landau gauge. Therefore, there is the following global $SU(2) \otimes SU(3)$ invariance:

$$e^{i\theta_W^a \mathcal{W}_a} \rightarrow U e^{i\theta_W^a \mathcal{W}_a} U^\dagger, \quad e^{i\theta_c^i C_i} \rightarrow \Gamma e^{i\theta_c^i C_i} \Gamma^\dagger \quad (6)$$

where $U \in SU(2)$ while $\Gamma \in SU(3)$. As a result we can represent equation (2) in the form:

$$N(\theta_Y, \theta_W, \theta_c, \theta_c') = \text{tr} [e^{i\theta_W \mathcal{W}_3} e^{i\theta_Y \mathcal{Y}} e^{i\theta_c C_8 + i\theta_c' C_3} \mathcal{N}] \quad (7)$$

where for the left-handed doublets of fermions

$$\mathcal{W}_3 = \frac{1}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad (8)$$

while for the colored quarks

$$C_8 = \frac{1}{3} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix}, \quad C_3 = \frac{1}{2} \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad (9)$$

The direct calculation gives

$$N(\theta_Y, \theta_W, \theta_c, \theta_c') = 2g \left(\cos \frac{\theta_Y}{2} - \cos \frac{\theta_W}{2} \right) \left(e^{i\theta_Y/6} \left(e^{\frac{i\theta_c}{3} + \frac{i\theta_c'}{2}} + e^{\frac{i\theta_c}{3} - \frac{i\theta_c'}{2}} + e^{-\frac{2i\theta_c}{3}} \right) + e^{-i\theta_Y/2} \right) \quad (10)$$

The particular case of this expression with $\theta_c = \theta_c' = 0$ was considered, for example, in [13, 26]. On the level of the angles θ_Y , θ_W , θ_c , θ_c' the Z_6 symmetry has the form:

$$\theta_Y \rightarrow \theta_Y + 2\pi N, \quad \theta_W \rightarrow \theta_W + 2\pi N, \quad \theta_c \rightarrow \theta_c + 2\pi N, \quad \theta_c' \rightarrow \theta_c' \quad (11)$$

(Notice that θ_W is defined as modulo 4π , θ_c is defined as modulo 6π , while θ_Y is defined as modulo 12π .) The generating function is robust to the deformations of the Green's function, if those deformations obey the SM symmetry.

The choice of parameters ($\theta_Y = 0$, $\theta_W = 2\pi$, $\theta_c = \theta_c' = 0$) and any other choice related to it by the Z_6 transformation of equation (11) gives the maximally possible value of the generating function:

$$N_{\max} = N(\theta_Y = 0, \theta_W = 2\pi, \theta_c = 0, \theta_c' = 0) = 16g. \quad (12)$$

This value guarantees that all 16g fermions of the SM are massless in the symmetric phase. Those maximal values (12) are formed by the discrete subgroup of the SM symmetry group [27] (that is related by the Z_6 transformation to the centers of $SU(3)_c$ and $SU(2)_L$):

$$N_{\max} = \text{tr} [K_Y \mathcal{N}] = 16g, \quad (13)$$

where we may take

$$K_Y = e^{i[2\pi(N+1)]_{\text{mod } 4\pi} \mathcal{W}_3} e^{i[2\pi N]_{\text{mod } 12\pi} \mathcal{Y}} e^{i[2\pi N]_{\text{mod } 6\pi} \mathcal{C}_8} \quad (14)$$

with any integer N . In particular, for $N = 3$ we get $K_Y = e^{6\pi i \mathcal{Y}}$ while for $N = 0$ we have $K_Y = e^{i2\pi \mathcal{W}_3}$. Various operators in equation (14) are related to the Z_6 transformation.

3. SM at low temperatures as the topological insulator with C, P, and T symmetries

The topology of the SM vacuum in the massive phase looks similar to that of the ground state of superfluid $^3\text{He-B}$, which in the noninteracting case is described by the integer valued topological invariant [24]

$$N_K = \frac{1}{24\pi^2} e_{\mu\nu\lambda} \text{tr} \left[K \int d^3p \mathcal{H}^{-1} \partial_{p_\mu} \mathcal{H} \mathcal{H}^{-1} \partial_{p_\nu} \mathcal{H} \mathcal{H}^{-1} \partial_{p_\lambda} \mathcal{H} \right]. \quad (15)$$

Here $\mathcal{H}(\mathbf{p}) = \mathcal{G}^{-1}(p_4 = 0)$ (while \mathcal{G} is the Green's function); the integral is over 3-momentum space; and K is the proper symmetry operation (it should either commute or anti-commute with \mathcal{H}). For the superfluid $^3\text{He-B}$ one has $N_K = 2$ for $K = \tau_2$ (the combination of time reversal and particle-hole symmetries [24]). The larger values of this invariant may be obtained by the extension of the model of $^3\text{He-B}$ to the multi-component fermionic models [28].

The expression for equation (15) is formally defined at zero temperature, $T = 0$. However the effects of the nontrivial topology on the physical properties of the systems can be measured at finite temperatures. For some effects the temperatures must be much smaller than the masses of the fermions existing in the given system, $T \ll m$, while for the others the limit $1/\tau(T) \ll m$ is enough, where $\tau(T)$ is the characteristic relaxation time. In the latter case it is not excluded that the topological invariant can be applicable even for $T > m$. Possibly, the definition of the topological invariant may be extended even further, but we do not discuss this possibility here.

The smallest Dirac mass in the SM is the mass of electron. The leading term in the temperature corrections to the corresponding self energy is the one loop expression proportional to the fine structure constant α . This term gives rise to the shift of the dispersion of the quasiparticles by the amount of the order of eT [60]. Therefore, the requirement $eT \ll m_e$ gives $T \ll 1$ MeV. An even greater restriction comes from the neutrino sector, where the thermal contribution to mass may be roughly given by the expression gT [61], where g is the $SU(2)$ or $U(1)_Y$ coupling constant. Assuming that the neutrino mass is about 1 eV, we arrive at the restriction $T \ll 1$ eV. This condition is satisfied, for example, by the present state of the universe with the temperature of the order of 10^{-4} eV.

The question arises, whether the vacuum of the SM in the massive phase is topologically trivial or not. If yes, what is the corresponding matrix K for the SM and what is the effect of interactions. The situation here is completely unclear. First of all, we may consider the approximation to the SM, in which the exchange by the W and Z bosons as well as the Higgs boson are neglected. Roughly, this may correspond to the description of processes at the energies much smaller than the electroweak scale ~ 100 GeV. Then in the SM at zero temperature $T = 0$ or at nonzero T with the proper restrictions such as $T \ll m$, where m is the smallest fermion mass in the phase with the spontaneously broken electroweak symmetry, the Green's function has the form

$$\mathcal{G}(p) = Z(-p^2) \frac{1}{\gamma^\mu p_\mu - M(-p^2)}, \quad (16)$$

where $p^2 = -\mathbf{p}^2 + \omega^2$, while Z and M are matrices. The fermion mass matrix m is given by the solution of equation

$$M(-m^2) = m$$

while the Dirac matrices may be chosen according to section 5.4 in [29]:

$$\gamma^0 = \tau_1, \quad \gamma = i\tau_2 \sigma, \quad \gamma_5 = -i\gamma^0 \gamma^1 \gamma^2 \gamma^3 = \tau_3. \quad (17)$$

This approximation is reasonable due to the smallness of the fine structure constant and large enough masses of W , Z , and the SM Higgs boson. Here it is the matrix $K = -i\gamma^5 \gamma^0 = \tau_2$, which commutes with the Green's function at $\omega = 0$. This is the matrix of the combination of CPT and P transformations⁸ that is at the same time the combination of C and T. As a result equation (15) determines the topological invariant. It may be calculated for the simplest system connected with the given one by a continuous transformation. Assuming that such a connection exists with the system of noninteracting massive Dirac fermions (that represent the constituents of the SM), we obtain

⁸ The unessential phase factor of this symmetry matrix is chosen in such a way that the expression of equation (15) gives the real value for the case of the noninteracting Dirac fermions.

$$N_K = 8g, \quad (18)$$

where g is the number of generations of the SM fermions⁹. In the notations in [30, 31] the invariant is $\nu = N_K/2$.

Presently the role of the interactions between the fermions is not completely clear. For example, even at zero temperature the strong $SU(3)$ interactions give rise to the transition between the system of the noninteracting quarks and the confining QCD. This may (but also may not) give rise to the value of the topological invariant associated with the CT symmetry that differs from the value calculated above. The answer depends on the possibility of continuously transforming the two-point fermion Green's function for the noninteracting massive fermions to the two-point Green's function of QCD with the strong interactions taken into account. Various approximations to QCD may provide different answers to this question. For example, the Nambu–Jona–Lasinio (NJL) approximation allows one to continuously connect the interacting and noninteracting Green's functions. The spectrum of the lightest resonances is described by the NJL model reasonably well. This allows us to suppose that in the low energy effective theory equation (18) gives the correct answer for the hadronic phase of the SM.

Formally the parity breaking interactions destroy the consideration of the topological stability based on the invariant $N_{K_{CT}}$. In practice, this invariant remains operative because of the smallness of the corrections. But the topological classification group may be reduced from Z to the smaller group. For example, the electroweak $SU(2)$ interactions assume that the vacua with the values of $N_{K_{CT}}$ of opposite signs (that correspond to the opposite values of the masses of all fermions) represent the same physical vacuum. This occurs because during the electroweak symmetry breaking the opposite values of masses appear as different versions of unitary gauge. In the complete theory with the $SU(2)$ interactions taken into account those states are related by global gauge transformation, and therefore, not only are continuously connected but represent the same physical vacuum. This reduces the topological classification to Z/Z_2 . The reduction may be more significant if the SM appears as a low energy approximation to a certain theory with the larger gauge group. For the recent discussion of the similar modification of the topological stability pattern in topological superconductors due to interactions see [30–38]. In practice the reduction of the topological classification means that various defects lose their topological stability. For example, let us consider the QCD sector of the SM with the two quarks (u and d). Let us also neglect the current masses of the quarks, There is the chiral $SU(2)$ symmetry which is broken spontaneously in the hadronic phase. As a result the constituent quark masses appear. The positive and negative values of the masses may appear in this way. One may naively suppose that this should lead to the formation of the topologically stable domain walls separating the regions with the opposite values of the constituent masses. But this is actually not so. The opposite values of the masses appear as the arbitrary choice of the sign of the condensate. Those choices are related to the element of the global chiral symmetry $SU(2)$. We may consider the version of the theory with the 2×2 complex-valued condensate field, and in this theory the state with positive mass is continuously connected by the symmetry transformation with the state with negative mass. (At the intermediate states the mass is undefined.) This means the reduction $Z \rightarrow Z/Z_2$ of the symmetry classification and this means that the topologically stable domain walls in the hadronic phase of QCD do not exist. In practice if such domain walls appear dynamically in the form of bubbles, then they decay with the emission of the $SU(2) \times U(1)$ gauge bosons.

4. Topological invariant protected by T-symmetry

4.1. Version of the SM with majorana masses of left-handed neutrinos

In section 3 we considered the approximation to the SM when parity remains unbroken. The question of the stability of vacuum was related to the topological invariant protected by CT. Interactions with the SM Higgs boson and with the W and Z bosons destroy the vacuum stability criteria based on the consideration of this invariant. At least, the interactions in the Higgs sector are strong. Although we may neglect this effect in some approximation, this is necessary to consider the other topological invariants. In order to consider such invariants we use the representation of the SM in terms of the Nambu–Gorkov spinors. This allows us to treat the particle–antiparticle transformation as a matrix. We assume in this section that weak CP breaking interactions do not affect the stability of vacuum. Therefore, we will use the topological invariant protected by T-symmetry.

In this subsection we consider the version of the SM with the left-handed massive neutrinos. First of all, let us discuss the situation, when the right-handed neutrinos remain massless, the Dirac masses of neutrinos are absent, and only the observed left-handed neutrinos are massive. For example, the type II neutrino seesaw may lead to such a pattern. Then the following mass term appears:

⁹ For the system of the noninteracting Dirac fermions with masses M_a the given topological invariant is given by $N_K = \sum_a \text{sign} M_a$.

$$L_\nu = -M \nu_L^A \nu_L^B \epsilon_{AB} + (\text{h.c.}) \quad (19)$$

(summation over the generations is implied).

Let us consider the situation, when the interacting system may be continuously deformed to the system without interactions in the lepton sector and with the Majorana masses of the left-handed neutrinos. Let us introduce the conventional definition of the Nambu–Gorkov spinor: $\mathcal{N}_L = (\nu_L^c, \nu_L)^T$ (where $\nu_L^c = i\sigma^2 \bar{\nu}_L$). In terms of this spinor the Lagrangian for one massive noninteracting left-handed neutrino may be written as follows:

$$L_L = \bar{\mathcal{N}}_L (\gamma^\mu p_\mu + M) \mathcal{N}_L \quad (20)$$

where

$$\bar{\mathcal{N}} = \mathcal{N}^T i\gamma^2 \gamma^0 \quad (21)$$

The time reversal transformation reads as:

$$\mathcal{N}_L \rightarrow \gamma^0 \gamma^5 \mathcal{N}_L \quad (22)$$

We choose the unessential phase factor in such a way that in this representation the matrix of the time reversal transformation is given by

$$\mathbf{K}_T = -i\gamma^0 \gamma^5 \quad (23)$$

(The corresponding topological invariant receives the form:

$$N_{K_T} = \frac{e_{ijk}}{48\pi^2} \text{tr} \left[\int_{\omega=0} d^3p \mathbf{K}_T G \partial_{p_i} G^{-1} G \partial_{p_j} G^{-1} G \partial_{p_k} G^{-1} \right]. \quad (24)$$

with

$$G^{-1} = \gamma^\mu p_\mu + M \quad (25)$$

We obtain:

$$\begin{aligned} N_{K_T} &= i \frac{e_{ijk}}{48\pi^2} \text{tr} \left[\int_{\omega=0} \frac{d^3p}{(\vec{p}^2 + M^2)^2} \gamma^0 \gamma^5 (+M) \gamma^i \gamma^j \gamma^k \right] \\ &= \frac{1}{8} \left(2 \times 4 \times \frac{1}{2} \right) = 1/2 \end{aligned} \quad (26)$$

If we have g generations of the left-handed neutrinos, the result is to be multiplied by g . As we will see below the quark sector with Dirac masses does not give the nonzero contribution to N_{K_T} . Therefore, assuming that the SM vacuum in the given phase is continuously connected with the vacuum of the version of the SM without interactions between leptons, we obtain the overall value of the invariant

$$N_{K_T} = g/2 \quad (27)$$

We suppose that this property takes place for the SM at vanishing temperature, pressure and chemical potentials.

4.2. Version of the SM with a type I neutrino seesaw

Let us remind the reader of the basics of the classical type I seesaw [41]. On the basis of $\mathcal{N}_{LR} = (\nu_L^c, \nu_R)^T$ (where $\nu_L^c = i\sigma^2 \bar{\nu}_L$) there is the mass matrix

$$\mathbf{M}_\nu = \begin{pmatrix} 0 & m \\ m & M \end{pmatrix} \quad (28)$$

The overall mass term is $\frac{1}{2} \mathcal{N}_{LR} \mathbf{M}_\nu \mathcal{N}_{LR} + (\text{h.c.})$, where (h.c.) means Hermitian conjugation which implies $\nu_{R,L} \rightarrow \bar{\nu}_{R,L}$ and vice versa. The product of the two-component spinors is defined as:

$$\mathcal{N}_{LR} \mathcal{N}_{LR} \equiv \mathcal{N}_{LR}^A \mathcal{N}_{LR}^B \epsilon_{AB}$$

For simplicity we assume that g Dirac masses m are equal to each other and g Majorana masses M are also equal. The diagonalization gives g heavy neutrinos with Majorana masses $M_{\text{heavy}} \approx +M$ and g light neutrinos with Majorana masses

$$-M_{\text{light}} \approx -m \frac{m}{M} \quad (29)$$

Notice that the signs of M_{heavy} and $-M_{\text{light}}$ are opposite. If we rewrite the mass term through the spinor ν_L rather than ν_L^c , then the sign of the mass becomes positive because

$$\begin{aligned} L_\nu &= -M_{\text{light}} [\nu_L^c]^T (i\sigma^2) \nu_L^c - M_{\text{light}} [\bar{\nu}_L^c]^T (-i\sigma^2) \bar{\nu}_L^c \\ &= M_{\text{light}} [\bar{\nu}_L]^T (-i\sigma^2) \bar{\nu}_L + M_{\text{light}} [\nu_L]^T (i\sigma^2) \nu_L \end{aligned} \quad (30)$$

This results in the trivial value of the corresponding momentum space topological invariant (see below). The assumptions that the mass of the right-handed neutrinos M is not smaller, than 1 TeV, and that the Dirac neutrino mass m is of the order of the electron mass m_e allow us to estimate $M_{\text{light}} \leq 0.25$ eV.

The topological invariant for the left-handed neutrino was calculated above and is given by equation (27). Now let us consider the right-handed neutrino. We define $\mathcal{N}_R = (\nu_R, \nu_R^c)^T$ (where $\nu_R^c = i\sigma^2 \bar{\nu}_R$). In terms of this spinor the lagrangian for the massive right-handed neutrino may be written as:

$$L_R = \bar{\mathcal{N}}_R (\gamma^\mu p_\mu + M) \mathcal{N}_R \quad (31)$$

Now the time reversal transformation reads as:

$$\mathcal{N}_R \rightarrow -\gamma^0 \gamma^5 \mathcal{N}_R \quad (32)$$

which means that in this representation

$$\mathbf{K}_T = i\gamma^0 \gamma^5 \quad (33)$$

The topological invariant is given by the same expression of equation (24). It gives:

$$N_{K_T} = -ig \frac{e_{ijk}}{48\pi^2} \text{tr} \left[\int_{\omega=0} \frac{d^3p}{(\vec{p}^2 + M^2)^2} \gamma^0 \gamma^5 (+M) \gamma^i \gamma^j \gamma^k \right] = -g/2 \quad (34)$$

where g is the number of generations. One can see that the system with the equal number of left-handed and right-handed neutrinos with Majorana masses of the same sign has the vanishing value of topological invariant.

4.3. Version of the SM with Dirac masses of neutrinos

In this subsection we discuss the case, where Majorana masses are absent, and follow the alternative definition of the Nambu–Gorkov spinors introduced in [39]. In appendix B we represent the corresponding definition. In terms of the corresponding Green's functions the topological invariant for the SM may be written as

$$N_K = \frac{e_{ijk}}{48\pi^2} \text{tr} \left[\int_{\omega=0} d^3p \mathbf{K} G \partial_{p_i} G^{-1} G \partial_{p_j} G^{-1} G \partial_{p_k} G^{-1} \right]. \quad (35)$$

where \mathbf{K} is the appropriate symmetry of the system while G is the Green's function being the vacuum average of the product of spinors Ψ of equation (43) of appendix B. For the definition of the two sets of gamma-matrices γ^μ and Γ^a also see the appendix B. First of all let us consider again the case of the P-invariant approximation to the SM, in which case we may use matrix of CT

$$\mathbf{K} = \mathbf{K}_{CT} = i\Gamma^4 \Gamma^5 \gamma^0$$

then if the Green's function of the system may be adiabatically deformed to that of the system with free massive Dirac fermions with the same mass M , then we can substitute into equation (35) the Green's function of the form:

$$G^{-1}(p) = i\gamma^2 \gamma^5 \gamma^0 \Gamma^4 \Gamma^2 \Gamma^5 [\gamma^\mu p_\mu + M\gamma^5 \Gamma^4] \quad (36)$$

This gives

$$N_{K_{CT}} = ig(N_c + 1) \frac{e_{ijk}}{48\pi^2} \text{tr} \left[\int_{\omega=0} \frac{d^3p}{(\vec{p}^2 + M^2)^2} \Gamma^4 \Gamma^5 \gamma^0 (\vec{\gamma} \vec{p} - M_0 \Gamma^4 \Gamma^5 \gamma^5) \gamma^i \gamma^j \gamma^k \right] = 2g(N_c + 1) \quad (37)$$

(where $N_c = 3$ is the number of colors) in accordance with the calculation of section 3.

If we take into account interactions that break parity, but neglect the complexness of the elements of the fermion mixing matrix, then the SM is CP-invariant. In this situation the topological invariant may be composed using the matrix

$$\mathbf{K}_T = -i\Gamma^4 \Gamma^2 \Gamma^5 \gamma^0$$

Let us again suppose that the two-point fermion Green's function of the SM is continuously connected to that of the noninteracting theory with all Dirac masses of the fermions equal to each other. In this case our topological invariant has the form

$$N_{K_T} = -ig(N_c + 1) \frac{e_{ijk}}{48\pi^2} \text{tr} \left[\int_{\omega=0} \frac{d^3p}{(\vec{p}^2 + M^2)^2} \Gamma^4 \Gamma^2 \Gamma^5 \gamma^0 (\vec{\gamma} \vec{p} - M_0 \Gamma^4 \Gamma^5 \gamma^5) \gamma^i \gamma^j \gamma^k \right] \sim \text{tr} \Gamma^2 \equiv 0 \quad (38)$$

Thus, one can see that the parity breaking interactions make the SM vacuum with Dirac masses of all fermions trivial.

5. Conclusions

In this paper we look at various phases of the SM of fundamental interactions (or its extensions) as at the systems similar to the topological materials. The symmetric phase (the temperature is above the temperature of the electroweak transition) represents the topological semimetal with the Fermi point, which is protected by the topological invariants generated by the functional equation (7). This functional equation depends on the angles θ_Y , θ_W^a , θ_c^i . It has been demonstrated that the Z_6 symmetry of the fermionic representations of the SM [25] manifests itself as the symmetry of this functional equation under the corresponding transformation of the mentioned angles in equation (3). Due to the Z_6 symmetry the maximal value of the topological invariant protecting the Fermi point may be constructed either of the hypercharge or of the generator \mathcal{W}_3 of $SU(2)_L$. It is given by equation (14). This maximal value is equal to the number of massless SM fermions. This is peculiar, although the formal definition of these topological invariants was given at zero temperature, this is the phase with high temperature, where they may be applied. This occurs because in this phase the mass scale disappears, and the relevant scale parameter is given by the temperature itself. It should be compared to the scale Λ , at which the SM transfers to its ultraviolet completion. We suppose that the topological invariants considered here protecting the Fermi point remain at work at least as long as $T \ll \Lambda$. Notice that such a scale may not be smaller than 1 TeV.

At the temperatures below the electroweak phase transition the SM (and its extensions) may exist in several phases, where the fermions are massive. Those phases resemble various phases of topological insulators and are characterized by the corresponding topological invariants in momentum space. At small enough temperature, pressure and chemical potentials the parity breaking interactions may be neglected, and the vacuum is topologically nontrivial being protected by the topological invariant of equation (35) with \mathbf{K} given by the matrix of CT transformation. Our definition of this invariant remains valid at least up to the temperatures much smaller than the neutrino masses, which are assumed to be of the order of eV . Although the strong interactions of the SM provide the transition between the system of noninteracting quarks and the quarks confined by the quark–gluon strings, we may suppose that the two-point quark Green’s function is continuously connected to that of the noninteracting quarks. At least this occurs in the NJL approximation to the QCD [56], which describes the light mesons reasonably well. See also the recent lattice data on the quark propagator in [59] and references therein, which indicate that the functions $Z(-p^2)$ and $M(-p^2)$ in equation (16) tend to finite nonzero values¹⁰ at $p \rightarrow 0$. Assuming the possibility of such continuous transformation, we come to the value of the topological invariant that protects the number of massive Dirac fermions.

If we take into account the parity broken interactions, then the topological classification based on the invariant protected by CT is reduced at least to Z/Z_2 . The further reduction is possible if the BSM unified model has the appropriate extended gauge symmetry. We deal with the topological insulator with T-symmetry if we neglect weak CP breaking interactions originated from the imaginary parts of the elements of the quark mixing matrix. Therefore, the stability of vacuum is protected also by the topological invariant of equation (35) with \mathbf{K} given by the matrix of the time reversal transformation. This expression remains operative at least for the temperatures $T \ll 1$ eV. We do not exclude the existence of the extension of its definition to the essentially higher temperatures. But this is beyond the scope of this paper. It appears that the value of this invariant for the SM with massive Dirac neutrinos is zero.

At the same time, the version of the SM with Majorana masses of left-handed neutrinos belongs to the topological class different from that of the SM with Dirac masses of the neutrinos. This means that the corresponding two systems cannot be connected continuously (at least, if we neglect the CP breaking interactions). For example, the Majorana masses of the left-handed neutrinos may appear as a result of the type II seesaw [49]. Considering the case of the type I seesaw, we come to the conclusion that the corresponding vacuum is topologically equivalent to the Dirac vacuum without Majorana masses, which follows from the existence of the Majorana masses of the right-handed neutrinos.

Acknowledgments

The part of the work of MAZ performed in Russia was supported by Russian Science Foundation Grant No 16-12-10059 (sections 2, 5, and appendix) while the part of the work done in France (the remaining part of the text) was supported by the Le Studium Institute of Advanced Studies. GEV acknowledges funding from the European Research Council (ERC) under the European Union’s Horizon 2020 Research and Innovation Program (Grant Agreement # 694248).

¹⁰ Though one cannot exclude considering these data that the quark function $Z(-p^2)$ tends to zero at $p \rightarrow 0$, in which case the singularity is encountered in the expression for the topological invariants. How to regularize such singularities if they do appear is a difficult question, and we omit this question in this paper.

Appendix A. Green's functions

We consider the two-point Green's function \mathcal{G} defined in a certain gauge corresponding to the gauge fixing condition $O[A] \rightarrow \min$.

$$\mathcal{G}_J^I(x, y) = \int D\bar{\psi}D\psi DA \exp \{-S[A, \psi]\} \exp \{-\lambda O[A] + \log \Delta_{FP}[A]\} \bar{\psi}_J(x) \psi^I(y), \quad (39)$$

Here the integral is over the fermionic fields of the SM while indices I, J enumerate the components of ψ . $S[A, \psi]$ is the action of the SM, A is the gauge field, the Faddeev–Popov determinant has the form:

$$\Delta_{FP}^{-1}[A] = \int dg \exp \{-\lambda O[A^g]\} \quad (40)$$

Here g is the gauge transformation and A^g is the transformed gauge field; it is implied that $\lambda \rightarrow \infty$ at the end. In our case the elements P of the gauge group G are unitary matrices. That is why $[\mathcal{G}, P] = 0$ means that $P^+ \mathcal{G} P = \mathcal{G}$.

$$\begin{aligned} [P^+]_L^I \mathcal{G}_J^I(x, y) P_I^K &= \int D\bar{\psi}D\psi DA \exp \{-S[A, P^+\psi]\} \\ &\quad \exp \{-\lambda O[A] + \log \Delta_{FP}[A]\} \bar{\psi}_L(x) \psi^K(y) \\ &= \int D\psi DA \exp \{-S[A, \psi]\} \\ &\quad \exp \{-\lambda O[P^+AP] + \log \Delta_{FP}[A]\} \bar{\psi}_L(x) \psi^K(y), \end{aligned} \quad (41)$$

From the last equation we obtain that $[\mathcal{G}, P] = 0$ for P from the center of G for any given gauge. At the same time when the functional $O[A]$ is invariant under the global gauge transformations, we also have $[\mathcal{G}, P] = 0$ for any $P \in G$. The particular case of such a gauge is the Landau gauge: $O[A] = \int \text{Tr} A^2 d^4x$. In this gauge any $P \in G$ commutes with the Green function. In the following we assume that this gauge is chosen.

Appendix B. Representation of the SM in terms of Nambu–Gorkov spinors

We adopt the notations proposed in [39]. Left-handed doublets and the right-handed doublets of quarks are denoted by L_K^a and R_K^a , where \mathbf{a} is the generation index while K is the color index. The left-handed doublets and the right-handed doublets of leptons are \mathcal{L}^a and \mathcal{R}^a , respectively. It will be useful to identify the lepton of each generation as the fourth component of colored quark. Then $L_{a,4}^a = \mathcal{L}_a^a$ and $R_{a,4}^a = \mathcal{R}_a^a$. So, later we consider the lepton number as the fourth color in the symmetric expressions. We define the analog of the Nambu–Gorkov spinor as:

$$\mathbf{L}_{aiU}^{\mathbf{a}A} = \begin{pmatrix} L_{ai}^{\mathbf{a}A} \\ \bar{L}_{c'i}^{\mathbf{a}B} \epsilon_{c'a} \epsilon^{BA} \end{pmatrix}, \quad \mathbf{R}_{aiU}^{\mathbf{a}A} = \begin{pmatrix} \bar{R}_{bi}^{\mathbf{a}B} \epsilon_{ba} \epsilon^{BA} \\ R_{a,i}^{\mathbf{a}A} \end{pmatrix},$$

where A is the usual spin index, U is the Nambu–Gorkov spin index ($U = 1, 2$ and $A = 1, 2$), i is the $SU(4)$ Pati–Salam color index (the lepton number is the fourth color), \mathbf{a} is the generation index, and a, b are the $SU(2)_L, SU(2)_R$ indices. Both $\mathbf{R}_{aiU}^{\mathbf{a}A}$ and $\mathbf{L}_{aiU}^{\mathbf{a}A}$ for the fixed values of a, i , and \mathbf{a} compose the four-component Dirac spinors $\mathbf{R}_{ai}^{\mathbf{a}}$ and $\mathbf{L}_{ai}^{\mathbf{a}}$. These spinors for the fixed value of a have $(N_c + 1) \times N_g = 12$ components. Both \mathbf{R} and \mathbf{L} belong to the fundamental representation of $U((N_c + 1) \times N_g)$, where $N_c = 3$ is the number of colors, $N_g = 3$ is the number of generations. Notice that \mathbf{L} and $\bar{\mathbf{L}}$ (\mathbf{R} and $\bar{\mathbf{R}}$) are not independent:

$$\bar{\mathbf{L}}_{ai}^{\mathbf{a}} = \epsilon_{ab} (\mathbf{L}_{bi}^{\mathbf{a}})^T i\gamma^2 \gamma^5 \gamma^0, \quad \bar{\mathbf{R}}_{ai}^{\mathbf{a}} = \epsilon_{ab} (\mathbf{R}_{bi}^{\mathbf{a}})^T i\gamma^2 \gamma^5 \gamma^0$$

Here the gamma-matrices act on the spinor space-time indices and are defined in chiral representation

$$\gamma^0 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \gamma^i = \begin{pmatrix} 0 & \sigma^i \\ -\sigma^i & 0 \end{pmatrix}, \quad i = 1, 2, 3, \quad \gamma^5 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad (42)$$

where σ^i is the space-time Pauli matrix.

Next, we arrange the Dirac spinors $\mathbf{L}_{ai}^{\mathbf{a}}, \mathbf{R}_{ai}^{\mathbf{a}}$ in the $SO(4)$ spinor Ψ :

$$\Psi_i^{\mathbf{a}} = \begin{pmatrix} \mathbf{L}_{ai}^{\mathbf{a}} \\ \mathbf{R}_{ai}^{\mathbf{a}} \end{pmatrix} \quad (43)$$

We introduce the Euclidean $SO(4)$ gamma-matrices Γ^a (in chiral representation). The action of the SM gauge fields $e^{i\theta} \in U(1)_Y \subset SU(2)_R, U^{(L)} \in SU(2)_L, U^{(R)} = \begin{pmatrix} e^{i\theta} & 0 \\ 0 & e^{-i\theta} \end{pmatrix} \in SU(2)_R$ and

$$V = \begin{pmatrix} Qe^{i\theta/3} & 0 \\ 0 & e^{-i\theta} \end{pmatrix} \in SU(4)_{\text{Pati-Salam}} \subset U(12) \text{ (where } Q \in SU(3)) \text{ on the given Majorana spinor is:}$$

$$\Psi_i^a \rightarrow \left(V_{ij} \frac{1 + \Gamma^5 \gamma^5}{2} + \bar{V}_{ij} \frac{1 - \Gamma^5 \gamma^5}{2} \right) \begin{pmatrix} U^{(L)} & 0 \\ 0 & U^{(R)} \end{pmatrix} \Psi_j^a$$

Thus $U^{(L)}$, $U^{(R)}$ realize the representation of $O(4) \simeq SU(2)_L \otimes SU(2)_R$ while V realizes the representation of the subgroup $SU(4)$ of $U(12)$. The action of the element $R \in U(12)$ of the latter group on the spinor Ψ_i^a is $\Psi_i^a \rightarrow \left(R_{ij}^{ab} \frac{1 + \Gamma^5 \gamma^5}{2} + \bar{R}_{ij}^{ab} \frac{1 - \Gamma^5 \gamma^5}{2} \right) \Psi_j^b$.

Again, Ψ and $\bar{\Psi}$ are not independent:

$$\bar{\Psi}_i^a = (\Psi_i^a)^T i\gamma^2 \gamma^5 \gamma^0 \Gamma^4 \Gamma^2 \Gamma^5$$

The gamma-matrices act on the internal $SU(2)_L$ and $SU(2)_R$ spinor indices rather than on the space-time indices, and are defined in chiral representation

$$\Gamma^4 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \Gamma^i = \begin{pmatrix} 0 & i\tau^i \\ -i\tau^i & 0 \end{pmatrix}, \quad i = 1, 2, 3, \quad \Gamma^5 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad (44)$$

Here τ^i are the basis elements of $SU(2)_{L,R}$ algebra. It is worth mentioning that matrix Γ^5 distinguishes the left-handed spinors from the right-handed spinors: $\Gamma^5 \mathbf{L} = \mathbf{L}$, and $\Gamma^5 \mathbf{R} = -\mathbf{R}$. In appendix C we list various symmetry operations that act on the spinors in terms of the Nambu–Gorkov spinor Ψ .

The partition function for the SM fermions in the presence of the SM gauge fields and the SM Higgs boson written in terms of spinor Ψ has the form: $Z = \int D\Psi e^{iS}$. The action $S = S_K + S_H$ contains two terms. The first one is the kinetic term

$$S_K = \frac{i}{2} \int d^4x (\bar{\Psi}_i^a \gamma^\mu \nabla_\mu \Psi_i^a) \quad (45)$$

One can check that this term being written in terms of the original SM fermions is reduced to the conventional SM fermion action (without mass term). Here ∇_μ is the covariant derivative that includes the gauge field of the model. The term S_H contains the interactions with the SM Higgs field. It appears in the form

$$\mathbf{H} = \sum_{K=1,2,3,4} \mathbf{h}_K \Gamma^K$$

where $\mathbf{h}^K \in \mathcal{R}$. Thus, the Higgs boson appears as the four-component real vector that is transformed under the action of $O(4) \simeq SU(2)_L \otimes SU(2)_R$. The following term would give equal masses to all fermions of the SM:

$$S_H = \frac{1}{2} \int d^4x (\bar{\Psi}_i^a \gamma^5 \Gamma^5 \mathbf{H} \Psi_i^a) \quad (46)$$

Using local $SU(2)_L \subset O(4) \approx SU(2) \otimes SU(2)$ transformation $L_{ai}^{aA} \rightarrow [U_{\mathbf{h}}]_{ab} L_{bi}^{aA}$ we may fix the unitary gauge, in which

$$\mathbf{H} = H\Gamma^4, \quad H = v + h \in \mathcal{R}, \quad (47)$$

(where h is the real-valued field of the 125 GeV Higgs boson while v is the condensate).

In order to obtain different masses for the fermions we need to introduce the matrix of the couplings between the Higgs field and the fermions. The corresponding term in the Lagrangian may be easily written in terms of the spinor Ψ , but we do not need this expression in our present consideration.

Below we represent the symmetries of space-time spinors in terms of the Nambu–Gorkov spinors introduced above.

1. CP transformation.

For the usual Dirac spinors ψ the CP transformation is:

$$\psi(t, \vec{r}) \rightarrow i\gamma^2 \bar{\psi}^T(t, -\vec{r})$$

In the left-right components we have:

$$\psi_L(t, \vec{r}) \rightarrow i\sigma^2 \bar{\psi}_L^T(t, -\vec{r})$$

and

$$\psi_R(t, \vec{r}) \rightarrow -i\sigma^2 \bar{\psi}_R^T(t, -\vec{r})$$

In terms of the spinors introduced above we have

$$\mathbf{L}_{ai}^a(t, \vec{r}) \rightarrow -\epsilon_{ab} \gamma^0 \mathbf{L}_{bi}^a(t, -\vec{r}), \quad \mathbf{R}_{ai}^a(t, \vec{r}) \rightarrow \epsilon_{ab} \gamma^0 \mathbf{R}_{bi}^a(t, -\vec{r})$$

and

$$\Psi_i^a(t, \vec{r}) \rightarrow \Gamma^4 \Gamma^2 \gamma^0 \Psi_i^a(t, -\vec{r})$$

2. Time reversal transformation.

In this paper we follow the definition of the time reversal transformation accepted in [40]. The time reversal transformation contains complex conjugation, which transforms the spinor ψ to its conjugate $\gamma^0 \bar{\psi}^T = \psi^*$ similar to the C transformation. At the same time the CPT transformation does not contain this conjugation, and being applied to usual spinors is composed of the multiplication by γ^5 and the inversion of time and space coordinates. According to the CPT theorem the T-transformation is equal to CP up to the overall inversion and the change in sign of the right-handed fermions. Therefore, the time reversal transformation results in

$$\Psi_i^a(t, \vec{r}) \rightarrow \Gamma^4 \Gamma^2 \Gamma^5 \gamma^0 \Psi_i^a(-t, \vec{r})$$

3. P transformation parity

$$\Psi_i^a(t, \vec{r}) \rightarrow \Gamma^4 \gamma^0 \Psi_i^a(t, -\vec{r})$$

4. Charge conjugation

Charge conjugation $C = CP \times P$:

$$\Psi_i^a(t, \vec{r}) \rightarrow \Gamma^2 \Psi_i^a(t, \vec{r})$$

5. CT transformation

$$\Psi_i^a(t, \vec{r}) \rightarrow \Gamma^4 \Gamma^5 \gamma^0 \Psi_i^a(-t, \vec{r})$$

References

- [1] Nielsen H B and Ninomiya M 1981 Absence of neutrinos on a lattice: I. Proof by homotopy theory *Nucl. Phys. B* **185** 20
- Nielsen H B and Ninomiya M 1981 Absence of neutrinos on a lattice: II. Intuitive homotopy proof *Nucl. Phys. B* **193** 173
- [2] Froggatt C D and Nielsen H B 1991 *Origin of Symmetry* (Singapore: World Scientific)
- [3] Grinevich P G and Volovik G E 1988 Topology of gap nodes in superfluid ^3He : π_4 homotopy group for ^3He -B disclination *J. Low Temp. Phys.* **72** 371
- [4] Hořava P 2005 Stability of Fermi surfaces and K -theory *Phys. Rev. Lett.* **95** 016405
- [5] Hasan M Z and Kane C L 2010 Topological insulators *Rev. Mod. Phys.* **82** 3045–67
- [6] Qi X-L and Zhang S-C 2011 Topological insulators and superconductors *Rev. Mod. Phys.* **83** 1057–110
- [7] Schnyder A P, Ryu S, Furusaki A and Ludwig A W W 2008 Classification of topological insulators and superconductors in three spatial dimensions *Phys. Rev. B* **78** 195125
- Schnyder A P, Ryu S, Furusaki A and Ludwig A W W 2009 Classification of topological insulators and superconductors *AIP Conf. Proc.* **1134** 10
- [8] Kitaev A 2009 Periodic table for topological insulators and superconductors *AIP Conf. Proc.* **1134** 22–30
- [9] Adler S 1969 Axial-vector vertex in spinor electrodynamics *Phys. Rev.* **177** 2426–38
- [10] Bell J S and Jackiw R 1969 A PCAC puzzle: $\pi_0 \rightarrow \gamma\gamma$ in the σ model *Nuovo Cim. A* **60** 47–61
- [11] Adler S L 2005 Anomalies to all orders *Fifty Years of Yang-Mills Theory* ed G 't Hooft (World Scientific) pp 187–228
- [12] Bevan T D C *et al* 1997 Momentogenesis by ^3He vortices: an experimental analog of primordial baryogenesis *Nature* **386** 689–92
- [13] Volovik G E 2003 *The Universe in a Helium Droplet* (Oxford: Clarendon Press)
- [14] Abrikosov A A and Beneslavskii S D 1970 Possible existence of substances intermediate between metals and dielectrics *ZhETF* **59** 1280–98
- Abrikosov A A and Beneslavskii S D 1971 *JETP* **32** 699
- [15] Soluyanov A A, Gresch D, Wang Z, Wu Q S, Troyer M, Dai X and Bernevig B A 2015 Type-II Weyl semimetals *Nature* **527** 495–8
- [16] Schnyder A P and Brydon P M R 2015 Topological surface states in nodal superconductors *J. Phys.: Condens. Matter* **27** 243201
- [17] Mizushima T, Tsutsumi Ya, Kawakami T, Sato M, Ichioka M and Machida K 2016 Symmetry protected topological superfluids and superconductors—From the basics to ^3He *J. Phys. Soc. Jpn.* **85** 022001
- [18] Bansil A, Lin H and Das T 2016 Colloquium: topological band theory *Rev. Mod. Phys.* **88** 021004
- [19] Yonezawa S 2016 Bulk topological superconductors *AAPPS Bulletin* **26** 3
- [20] Volovik G E and Zubkov M A On symmetry and topological origin of Weyl particles arXiv:1312.1267
- [21] Volovik G E and Zubkov M A 2014 Emergent Weyl spinors in multi-fermion systems *Nucl. Phys. B* **881** 514–38
- [22] Volovik G E and Zubkov M A 2014 Emergent Weyl fermions and the origin of $i = \sqrt{-1}$ in quantum mechanics *Pis'ma ZhETF* **99** 552–7
- Volovik G E and Zubkov M A 2014 *JETP Lett.* **99** 481–6
- [23] Zubkov M A 2007 The observability of $Z(6)$ symmetry in the standard model *Phys. Lett. B* **649** 91
- Zubkov M A 2007 *Phys. Lett. B* **655** 309 (erratum)

- [24] Volovik G E 2009 Fermion zero modes at the boundary of superfluid $^3\text{He-B}$ *Pis'ma ZhETF* **90** 440–2
 Volovik G E 2009 *JETP Lett.* **90** 398–401
 Volovik G E 2009 Topological invariant for superfluid $^3\text{He-B}$ and quantum phase transitions *Pis'ma ZhETF* **90** 639–43
 Volovik G E 2009 *JETP Lett.* **90** 587–91
 Volovik G E 2010 Topological superfluid $^3\text{He-B}$ in magnetic field and Ising variable *Pis'ma ZhETF* **91** 215–9
 Volovik G E 2010 *JETP Lett.* **91** 201–5
- [25] Bakker B L G, Veselov A I and Zubkov M A 2004 Internal structure of discretized Weinberg–Salam model *Phys. Lett. B* **583** 379
- [26] Volovik G E 2000 Momentum-space topology of standard model *J. Low Temp. Phys.* **119** 241–7
- [27] Volovik G E 2010 Topological invariants for standard model: from semi-metal to topological insulator *Pis'ma ZhETF* **91** 61–7
 Volovik G E 2010 *JETP Lett.* **91** 55–61
- [28] Yang W, Li Y and Wu C 2016 Topological septet pairing with spin-3/2 fermions: high-partial-wave channel counterpart of the $^3\text{He-B}$ phase *Phys. Rev. Lett.* **117** 075301
- [29] Weinberg S 1996 *The Quantum Theory of Fields* (Cambridge: Cambridge University Press)
- [30] Metlitski M A, Fidkowski L, Chen X and Vishwanath A Interaction effects on 3D topological superconductors: surface topological order from vortex condensation, the 16 fold way and fermionic Kramers doublets arXiv:1406.3032
- [31] Ryu S 2015 Interacting topological phases and quantum anomalies *Phys. Scr.* **164** 014009
- [32] Tachikawa Y and Yonekura K Gauge interactions and topological phases of matter arXiv:1604.06184
- [33] Witten E Fermion path integrals and topological phases arXiv:1508.04715
- [34] Yonekura K Dai–Freed theorem and topological phases of matter arXiv:1607.01873
- [35] Hsieh C-T, Cho G Y and Ryu S 2016 Global anomalies on the surface of fermionic symmetry-protected topological phases in (3+1) dimensions *Phys. Rev. B* **93** 075135
- [36] Wang C and Senthil T 2014 Interacting fermionic topological insulators/superconductors in three dimensions *Phys. Rev. B* **89** 195124
- [37] Kapustin A, Thorngren R, Turzillo A and Wang Z 2015 Fermionic symmetry protected topological phases and cobordisms *J. High Energy Phys.* **JHEP12(2015)052**
- [38] Kitaev A Homotopy-theoretic approach to SPT phases in action: Z_{16} classification of three-dimensional superconductors (<http://www.ipam.ucla.edu/abstract/?tid=12389&pcode=STQ2015>)
- [39] Zubkov M A 2014 Modified model of top quark condensation *Phys. Rev. D* **90** 057501
- [40] Berestetskii V B, Lifshitz E M and Pitaevskii L P 1982 Quantum electrodynamics *Course of Theoretical Physics* vol 4 2nd edn (Oxford: Pergamon Press)
- [41] Hernandez P *Neutrino physics, CERN Yellow Report* CERN-2010-001 229–78 arXiv:1010.4131
- [42] So H 1985 Induced topological invariants by lattice fermions in odd dimensions *Prog. Theor. Phys.* **74** 585–93
- [43] Ishikawa K and Matsuyama T 1986 Magnetic field induced multi component QED in three-dimensions and quantum Hall effect *Z. Phys. C* **33** 41–5
- [44] Kaplan D B 1992 Method for simulating chiral fermions on the lattice *Phys. Lett. B* **288** 342–7
- [45] Golterman M F L, Jansen K and Kaplan D B 1993 Chern–Simons currents and chiral fermions on the lattice *Phys. Lett. B* **301** 219–23
- [46] Creutz M 2008 Four-dimensional graphene and chiral fermions *J. High Energy Phys.* **JHEP04(2008)017**
- [47] Kaplan D B and Sun S 2012 Spacetime as a topological insulator: mechanism for the origin of the fermion generations *Phys. Rev. Lett.* **108** 181807
- [48] Creutz M 2011 Confinement, chiral symmetry, and the lattice *Acta Phys. Slovaca* **61** 1–127
- [49] Grimus W, Lavoura L and Radovic B 2009 Type II seesaw mechanism for Higgs doublets and the scale of new physics *Phys. Lett. B* **674** 117
- [50] Zubkov M A and Volovik G E 2012 Momentum space topological invariants for the 4D relativistic vacua with mass gap *Nucl. Phys. B* **860** 295–309
- [51] Zubkov M A Momentum space topology in the lattice gauge theory arXiv:1210.1989
- [52] Alford M, Rajagopal K and Wilczek F 1999 Color-flavor locking and chiral symmetry breaking in high density QCD *Nucl. Phys. B* **537** 443–58
- [53] Buballa M 2005 NJL-model analysis of dense quark matter *Phys. Rept.* **407** 205–376
- [54] Anglani R, Mannarell M and Ruggieri M 2011 Collective modes in the color flavor-locked phase *New J. Phys.* **13** 055002
- [55] Eto M, Nitta M and Yamamoto N 2010 Instabilities of non-Abelian vortices in dense QCD *Phys. Rev. Lett.* **104** 161601
 Eto M and Nitta M 2009 Color magnetic flux tubes in dense QCD *Phys. Rev. D* **80** 125007
- [56] Volkov M K and Radzhabov A 2006 The Nambu–Jona–Lasinio model and its development *Phys. Usp.* **49** 551–61
- [57] Essin A M and Gurarie V 2011 Bulk-boundary correspondence of topological insulators from their Green’s functions *Phys. Rev. B* **84** 125132
- [58] Zubkov M A 2016 Wigner transformation, momentum space topology, and anomalous transport *Annals Phys.* **373** 298
- [59] Oliveira O, Kizilersu A, Silva P J, Skullerud J I, Sternbeck A and Williams A G Lattice Landau gauge quark propagator and the quark-gluon vertex arXiv:1605.09632 [hep-lat]
- [60] Bellac M L 1996 Thermal field theory *Cambridge Monographs on Mathematical Physics* (Cambridge University Press)
- [61] Ghiglieri J and Laine M 2016 Neutrino dynamics below the electroweak crossover *JCAP* **JCAP07(2016)015**